

PART B
SOLAR - GEOPHYSICAL DATA

ISSUED
FEBRUARY 1960

U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS
CENTRAL RADIO PROPAGATION LABORATORY
BOULDER, COLORADO

SOLAR - GEOPHYSICAL DATA

CONTENTS

INTRODUCTION

I DAILY SOLAR INDICES

- (a) Relative Sunspot Numbers and 2800 Mc Solar Flux
- (b) Graph of Sunspot Cycle

II SOLAR CENTERS OF ACTIVITY

- (a) Calcium Plage and Sunspot Regions
- (b) Coronal Line Emission Indices - January 1960

III SOLAR FLARES

- (a-c) Optical Observations - January 1960
- (d) Flare Patrol Observations - December 1959
- (e,f) Subflares - December 1959
- (g-k) Optical Observations - October 1959
- (l) Flare Patrol Observations - October 1959
- (m) Ionospheric Effects (SWF) - December 1959
- (n) Ionospheric Effects (SEA-SCNA-Bursts) - September 1959

IV SOLAR RADIO WAVES

- (a) 2800 Mc -- Outstanding Occurrences (Ottawa) January 1960
- (b) 167 Mc -- Outstanding Occurrences (Boulder) January 1960
- (c) 169 Mc -- Outstanding Occurrences (Nançay) January 1960

V GEOMAGNETIC ACTIVITY INDICES

- (a) C, Kp, Ap, and Selected Quiet and Disturbed Days
- (b) Charts of Kp by Solar Rotations

VI RADIO PROPAGATION QUALITY INDICES

North Atlantic:


- (a) CRPL Quality Figures and Forecasts
- (b) Graphs Comparing Forecast and Observed Quality
- (c,d) Graphs of Useful Frequency Ranges

North Pacific:

- (e) CRPL Quality Figures and Forecasts
- (f) Graphs Comparing Forecast and Observed Quality

VII ALERT PERIODS AND SPECIAL WORLD INTERVALS

- (a) IGC 1959 Alerts and SWI



SOLAR - GEOPHYSICAL DATA

INTRODUCTION

This monthly report series is intended to keep research workers abreast of the major particulars of solar activity and the associated ionospheric, radio propagation and other geophysical effects. It is made possible through the cooperation of many observatories, laboratories and agencies as recorded in the detailed description of the tables and graphs which follows. The report is prepared in the Radio Warning Services Section, edited by Miss J.V. Lincoln.

I DAILY SOLAR INDICES

Relative Sunspot Numbers -- The table includes (1) the daily American relative sunspot numbers, R_A , as compiled by the Solar Division of the American Association of Variable Star Observers, and (2) the provisional daily Zürich relative sunspot numbers, R_Z , as communicated by the Swiss Federal Observatory. Because of the time required to collect and reduce the observations, R_A will normally appear one month later than R_Z .

The relative sunspot number is an index of the activity of the entire visible disk. It is determined each day without reference to preceding days. Each isolated cluster of sunspots is termed a sunspot group and it may consist of one or a large number of distinct spots whose size can range from 10 or more square degrees of the solar surface down to the limit of resolution (e.g. $1/8$ square degrees). The relative sunspot number is defined as $R=K(10g+s)$, where g is the number of sunspot groups and s is the total number of distinct spots. The scale factor K (usually less than unity) depends on the observer and is intended to effect the conversion to the scale originated by Wolf. The observations for sunspot numbers are made by a rather small group of extraordinarily faithful observers, many of them amateurs, each with many years of experience. The counts are made visually with small, suitably protected telescopes.

Final values of R_Z appear in the IAU Quarterly Bulletin on Solar Activity, the Journal of Geophysical Research, these reports, and elsewhere. They usually differ slightly from the provisional values. The American numbers, R_A , are not revised.

Solar Flux Values, 2800 Mc -- The table also lists the daily values of solar flux at 2800 Mc recorded in watts/ M^2 /cycle/second bandwidth ($\times 10^{-22}$) in two polarizations by the National Research Council at Ottawa, Canada. These solar radio noise indices are being published in accordance with CCIR Report 25 that a basic solar index for ionospheric propagation should be measured objectively and "preferably refer to a property of the sun such as radiation flux which has direct physical relationship to the ionosphere."

Graph of Sunspot Cycle -- The graph illustrates the recent trend of Cycle 19 of the 11-year sunspot cycle and some predictions of the future level of activity. The customary "12-month" smoothed index, \bar{R} , is used throughout, the data being final R_Z numbers except for the current year. Predictions shown are those made for one year after the latest available datum by the method of A. G. McNish and J. V. Lincoln (Trans. Am. Geophys. Union, 30, 673-685, 1949) modified by the use of regression coefficients and mean cycle values recomputed for Cycles 8 through 18. Cycle 19 began April 1954, when the minimum \bar{R} of 3.4 was reached.

II SOLAR CENTERS OF ACTIVITY

Calcium Plage and Sunspot Regions -- The table gives particulars of the centers of activity visible on the solar disk during the preceding month. These are based on estimates made and reported on the day of observation and are therefore of limited reliability.

The table gives the heliographic coordinates of each center (taken as the calcium plage unless two or more significantly and individually active sunspot groups are included in an extended plage) in terms of the Greenwich date of passage of the sun's central meridian (CMP) and the latitude; the serial number of the plage as assigned by McMath-Hulbert Observatory; the serial number of the center in the previous solar rotation, if it is a persisting region; particulars of the plage at CMP: area, central intensity; a summary of the development of the plage during the current transit of the disk, where b = born on disk, ℓ = passed to or from invisible hemisphere, d = died on disk, and / = increasing, - = stable, \ = decreasing; and age in solar rotations; particulars of the associated sunspot group, if any, at CMP: area and spot count and the summary of development during the current disk transit, similar to the above. The unit of area is a millionth of the area of a solar hemisphere; the central intensity of calcium plages is roughly estimated on a scale of 1 = faint to 5 = very bright.

Calcium plage data are available through the cooperation of the McMath-Hulbert Observatory of the University of Michigan. The sunspot data are compiled from reports from the U. S. Naval Observatory, and from reports from Europe and Japan received through the daily Ursigram messages.

Coronal Line Emission Indices -- In the table are summarized solar coronal emission intensity indices for the green (Fe XIV at $\lambda 5303$) and red (Fe X at $\lambda 6374$) coronal lines. The indices are based on measurements made at 5° intervals around the periphery of the solar disk by the High Altitude Observatory at Climax, Colorado, and by Harvard University observers at Sacramento Peak (The USAF Upper Air Research Observatory at Sunspot, New Mexico, under contract AF 19(604)-146). The measurements are expressed as the number of millionths of

an Angstrom of the continuum of the center of the solar disk (at the same wavelength as the line) that would contain the same energy as the observed coronal line. The indices have the following meanings:

G_6 = mean of six highest line intensities in quadrant for $\lambda 5303$.

R_6 = same for $\lambda 6374$.

G_1 = highest value of intensity in quadrant, for $\lambda 5303$.

R_1 = same for $\lambda 6374$.

The dates given in the table correspond to the approximate time of CMP of the longitude zone represented by the indices. The actual observations were made for the North East and South East quadrants 7 days before; for the South West and North West quadrants 7 days after the CMP date given.

To obtain rough measures of the integrated emission of the entire solar disk in either of the lines, assuming the coronal changes to be small in a half solar rotation, it is satisfactory to perform the following type of summation given in example for 15 October:

$$\left(\text{MEAN DISK EMISSION} \right)_{15 \text{ OCT}} \text{ IN } \lambda 5303 = \frac{1}{N} \left[\sum_{15 \text{ OCT}}^{22 \text{ OCT}} \left\{ (G_6)_{\text{NE}} + (G_6)_{\text{SE}} \right\} + \sum_{8 \text{ OCT}}^{14 \text{ OCT}} \left\{ (G_6)_{\text{SW}} + (G_6)_{\text{NW}} \right\} \right]$$

where N is the number of indices entering the summation.

Such integrated disk indices as well as integrated whole-sun indices are computed for each day and are published quarterly in the "Solar Activity Summary" issued by the High Altitude Observatory at Boulder, Colorado. In the same reports are given maps of the intensity distribution of coronal emission derived from all available Climax and Sacramento Peak observations, as well as other information on solar activity, such as maps made from daily limb prominence surveys in $H\alpha$ and notes regarding the history of active regions on the solar disk.

Preliminary summaries of solar activity, prepared on a fast schedule, are issued Friday of each week from High Altitude Observatory in conjunction with CRPL and include solar activity through the preceding day. These are useful to groups needing information on the current status of activity on the visible solar disk, but are not recommended for research uses unless such a prompt schedule of reporting is essential. The same information is included in the subsequent quarterly reports, with extensive additions, corrections and evaluations.

III SOLAR FLARES

Optical Observations -- The table presents the preliminary record of solar flares as reported to the CRPL on a rapid schedule at the sacrifice of detailed accuracy. Definitive and complete data are published later in the Quarterly Bulletin on Solar Activity, I.A.U., in various observatory publications and elsewhere. The present listing serves to identify and roughly describe the phenomena observed.

Reporting directly to the CRPL are the following observatories: McMath-Hulbert, Wendelstein, Sacramento Peak, Mitaka and Swedish Astrophysical Station on Capri. The remainder report through the URSIgram centers or are available through the IGY World Data Center for Solar Activity in Boulder. Observations are in the light of the center of the H-alpha line unless noted otherwise. The reports from Sacramento Peak, New Mexico (communicated to CRPL by the High Altitude Observatory at Boulder) are from observations at the USAF Upper Air Research Observatory at Sunspot, New Mexico, by Harvard University observers, under contract AF 19(604)-4961.

For each flare are listed the reporting observatory, the date, beginning and ending times, time of maximum phase, the heliographic coordinates in degrees, McMath serial number of the region, duration, the flare importance on the IAU scale of 1- to 3+, observing conditions where 1 means poor, 2 fair and 3 good, time of measurement for tabulated width of H α or tabulated area, measured (i.e. projected) maximum area in square degrees, corrected maximum area in square degrees which equals measured area times secant h where h is the heliocentric angle, maximum effective line-width in H α expressed in Angstroms, and maximum intensity of H α expressed in per cent of the continuous spectrum. The following symbols are used in the table:

D = Greater than	F = Approximately
E = Less than	ϵ = Plus

A final column lists provisionally the occurrence of simultaneous ionospheric effects as observed on selected field-strength recordings of distant high-frequency radio transmissions; a more nearly definitive list of these ionospheric effects, including particulars, appears in these reports after the lapse of a month (see below). All times are Universal Time (UT or GCT). Subflares (importance 1-) are listed by date, time of beginning and their heliographic coordinates. A graph presents intervals for which there were no patrols for flare observations from the observatories whose complete data are published in the table.

Ionospheric Effects -- SID, sudden ionospheric disturbances (and GID--gradual ionospheric disturbances) may be detected in a number of ways: short wave fadeouts (SWF), enhancement of low frequency atmospherics (SEA), increases in cosmic absorption (SCNA), and so forth.

A table lists SWF events that have been recognized on field-strength recordings of distant high-frequency radio transmissions. Under a coordinated program, the staffs at the following ionospheric sounding stations contribute reports that are screened and synthesized at CRPL-Boulder: Puerto Rico, Ft. Belvoir, Va., and Anchorage, Alaska (CRPL Stations: PR, BE, AN); Huancayo, Peru (CRPL-Associated Laboratory: HU); and Ft. Monmouth, N.J., White Sands, N. Mex., Adak, Alaska, and Okinawa (U.S. Signal Corps Stations: FM, WS, AD, OK). McMath-Hulbert Observatory (MC) also contributes such reports. In addition, reports are volunteered by RCA Communications Inc., Marconi Wireless, Netherlands Postal and Telecommunications Services, Swedish Telecommunications, and others; these usually specify times of SWF and the radio paths involved. Through the URSIgrams, reports are available from still other stations as given monthly in the footnotes.

In the coordinated program, the abnormal fades of field strength not obviously ascribable to other causes, are described as short wave fadeouts with the following further classification:

- S-SWF: sudden drop-out and gradual recovery
- Slow S-SWF: drop-out taking 5 to 15 minutes and gradual recovery
- G-SWF: gradual disturbance; fade irregular in either drop-out or recovery or both.

When there is agreement among the various reporting stations on the time (UT) of an event, it is accepted as a widespread phenomenon and listed in the table.

The degree of confidence in identifying the event, a subjective estimate, is reported by the stations and this is summarized in an index of certainty that the event is widespread, ranging from 1 (possible) to 5 (definite). The times given in the table for the event are from the report of a station (underlined in table) that identified it with high confidence. The criteria for the subjective importance rating assigned by each station on a scale of 1- to 3+ include amplitude of the fade, duration and confidence; greater consideration is given to reports on paths near the subsolar point in arriving at the summary importance rating given in the table.

Note: The tables of SID observed at Washington included in CRPL F-reports prior to F-135 were restricted to events classed here as S-SWF.

A second table lists sudden ionospheric disturbances which have been recognized on recorders for detecting cosmic absorption at about 18 Mc (SCNA) or on recorders for detecting enhancements of low frequency atmospheric at about 27 kc (SEA) together with solar radio bursts at 18 Mc as identified on the SCNA records.

Reports are received either directly or through the IGY World Data Center for Solar Activity at the High Altitude Observatory, Boulder, Colo. The following observatories report SCNA: Rensselaer Polytechnic Institute Observatory, Grafton, N.Y. (RE); McMath-Hulbert

Observatory (MC); Sacramento Peak, N.Mex. (SP); High Altitude Observatory, Boulder, Colo. (BO); University of Hawaii, Makapuu Pt., Hawaii (HA); and the Royal Observatory Edinburgh (ED). All of these except the Royal Observatory Edinburgh also report solar noise bursts observed at 18 Mc. The SEA reports come from the following: Department of Terrestrial Magnetism, Carnegie Institution of Washington, Station at Derwood, Md. (DE); Dunsink Observatory, Ireland (DU); Royal Observatory Edinburgh (ED); three stations operated by the Netherlands PTT at Hollandia, Dutch West Indies (HO), Nederhorst den Berg, Netherland (NE), and Paramaribo, New Guinea (PA); Panska Ves Observatory near Prague, Czech. (PU); High Altitude Observatory, Boulder, Colo. (BO); Sacramento Peak, N.Mex. (SP); McMath-Hulbert Observatory (MC); University of Hawaii (HA); Neustrelitz (NU); Kuhlungsborn (KU); and a group of American Association of Variable Star Observers located at Brooklyn, N.Y. (A1), Pittsburgh, Pa. (A2), Paterson, N.J. (A3), Powell, Ohio (A4), Ramsey, N.J. (A5), Oshkosh, Wis. (A6), China Lake, Calif. (A7), and Manhattan, Kansas (A8).

These reports are coordinated at CRPL-Boulder. When there is agreement among the various reporting stations on the time (UT) of an event, it is accepted as a widespread phenomenon and listed in the table. Some phenomena are listed, if noted at only one location, if there has been a flare or another type of flare-associated effect reported for that time.

In the table under the type of event the importance of the event is given on a scale of 1 minus to 3 plus. Next there is the index of widespread certainty ranging from 1 (possible) to 5 (definite). The time of beginning, maximum and end of the event in UT is given as reported by the station underlined in the group of observing stations. If the event is an SCNA, a percent absorption figure is given. This absorption is calculated by

$$\text{SCNA \%} = \frac{I_n - I_f}{I_n} \times 100$$

where I_n = noise diode current required to give a recorder deflection equal to that which would have occurred in the absence of a flare, i.e. a value extrapolated from cosmic noise level trend before and after a flare. The previous day's record may be considered if necessary.

and I_f = noise diode current required to give a recorder deflection equal to the level at the time of maximum absorption.

IV SOLAR RADIO WAVES

2800 Mc Observations

The data on solar radio wave events made in Ottawa, Canada by the Radio and Electrical Engineering Division of the National Research Council (A. E. Covington) at 2800 Mc (10-cm emission) are presented. Near local noon (about 1700 UT) the sensitivity of the radiometer is determined and a mean flux for the whole day calculated. These values are given in a tabular form (see table I-1) in units of 10^{-22} watts/ $M^2/c/s$. Burst phenomena are measured above this level and are given in terms especially suitable for the variations observed on this frequency. The basis for the classifications is described by Covington - J.R. Astro. Soc. Can. 45, 49, 1951 and Dodson, Hedeman and Covington, Ap. J. 119, 541, 1954. A modification in terminology with a view to simplification has been introduced and consists essentially of the omission of the descriptive word "Single" from the "Single-Simple" and "Single-Complex" classes; in designating the "Single", "Single-Simple" and "Rise and Fall" bursts into a single classification designated as "Simple Bursts" with an appropriate type number; in the addition of the letter "f" to indicate that the burst deviates from the basic pattern by the presence of one or more small fluctuations in intensity; and by the addition of the letter "A" to indicate that the event has another smaller duration event superimposed upon it.

Simple Burst

Any single burst which rises to one maximum and then decreases to the pre-burst level.

1 - Simple 1 -- Simple burst, type 1 (formerly "single"). Bursts of intensity less than 7 1/2 flux units and duration less than 7 1/2 minutes.

2 - Simple 2 -- Simple burst, type 2 (formerly "single-simple"). Bursts of impulsive nature with intensity greater than 7 1/2 flux units.

3 - Simple 3 -- Simple burst, type 3 (formerly "rise and fall"). Bursts of moderate intensity with duration greater than 7 1/2 minutes.

4 - Post-burst increase -- Postburst level is greater than the preburst level. The gradual return to normal flux may require as long as several hours.

5 - Absorption following burst (negative post).

6 - Complex -- (formerly "single-complex"). A single burst which shows two or more comparable maxima before the activity has declined to zero.

7 - Period of irregular activity or fluctuations -- Series of overlapping bursts of moderate intensity and duration.

8 - Group -- Series of single isolated bursts occurring in succession with intensity between the events equal to the level before and after the group.

9 - Precursor -- A small increase of intensity occurring before a larger increase.

Great Burst


Infrequently occurring bursts of great intensity, often of complicated structure.

Letter "A"

Indicates that this event has another event superimposed upon it.

Letter "f"

Indicates that the basic form of the event is modified by secondary fluctuations.



200 Mc Observations

Data on solar radio emission on 200 Mc recorded by the University of Hawaii (I. Miyake) at Makapuu Pt., Hawaii, are presented. The outstanding occurrences are reported as described under 170 Mc Observations with the exception that no intensity measurements are given.

170 Mc Observations

Data on solar radio emission at the nominal frequency of 170 Mc recorded at the Gunbarrel Hill (Boulder) station of the National Bureau of Standards (C.G. Little) are presented. The half width of the antenna lobe is appreciably greater than the solar disk. Polarization is not determined, but the dipole is oriented E-W. All times are in Universal Time (UT or GCT). Observations are interrupted during the period from 26 to 29 minutes after each hour for calibrations.

Beginning January 1, 1959 the method of reducing the records has been changed. The 3-hourly and daily flux density and variability are no longer determined. The outstanding occurrences are reported. However, instead of giving the intensity to the nearest unit of 10^{-22} watts meter $^{-2}$ (c/s) $^{-1}$, a scale of 1 to 3 is now used where for the estimate of smoothed maximum flux:

- 1 signifies $<100 \times 10^{-22}$ $\text{wm}^{-2}(\text{c/s})^{-1}$
- 2 signifies $>100 <1000 \times 10^{-22}$ $\text{wm}^{-2}(\text{c/s})^{-1}$
- 3 signifies $>1000 \times 10^{-22}$ $\text{wm}^{-2}(\text{c/s})^{-1}$.

Starting and maximum times are read to the nearest 1/10 minute if they are very definite and otherwise to the nearest minute. If the duration is less than five minutes, it is given to the nearest 1/10 minute; otherwise to the nearest minute. The following qualifying symbols are used:

- E = Event in progress before observations began.
- D = Event continues after observations cease.
- I = Event apparently continued during an interruption of the observations. The period of the interruption may be given in the remarks.
- S = Measurement may be influenced by interference or atmospherics.

The types of the outstanding occurrences follow the classification described by Dodson, Hedeman and Owren (Ap J. 118, 169, 1953), in which the types are identified by numbers which describe the character of the trace, but not the magnitude of the event, as follows:

0 - Rise in base level -- A temporary increase in the continuum with duration of the order of tens of minutes to an hour.

1 - Series of bursts -- Bursts or groups of bursts, occurring intermittently over an interval of time of the order of minutes or hours. Such series of bursts are assigned as distinctive events only when they occur on a smooth record or show as a distinct change in the activity.

2 - Groups of bursts -- A cluster of bursts occurring in an interval of time of the order of minutes.

3 - Minor burst -- A burst of moderate or small amplitude, and duration of the order of one or two minutes.

4 - Minor burst and second part -- A double rise in flux in which the early rise is a minor burst.


6 - Noise storm -- A temporary increase in radiation characterized by numerous closely spaced bursts, by an increase in the continuum, or by both. Duration is of the order of hours or days.

7 - Noise storm begins -- The onset of a noise storm occurs at some time during the observing period.


8 - Major burst -- An outburst, or other burst of large amplitude and more than average duration. A major burst is usually complex, with a duration of the order of one to ten minutes.

9A, 9B, or 9 -- Major burst and second part or large event without distinct first and second parts -- If there is a double rise in flux, the first part, a major burst, is listed as 9A and the second part as 9B. The second part may consist of a rise in base level, a group or series of bursts, a noise storm. A major increase in flux with duration greater than ten minutes but without distinct first and second parts, is listed simply as 9.


0-RISE IN BASE LEVEL




1-SERIES




2 - GROUP




3 - MINOR




4 - MINOR+



7-ONSET OF NOISE STORM





Note: In the present table, the type classifications 0 and 1 are not used; they have been included above only for information.

169 Mc Interferometric Observations

The 169 Mc interferometric observations are recorded around local noon at Nançay (Cher), France, (N47°23', E8^m47^s) the field station of the Meudon Observatory.

The main lobes are parallel to the meridian plane: the half-power width is 3.8 minutes in the East-West direction and much larger than the solar diameter in the North-South direction. The main lobes are about 2° apart (Ann. Astrophys. 20, 155, 1957). The records give the strip intensity distribution from the center of the disk to 30° to the West and East.

These daily distributions are plotted on the same chart giving diagrams of evolution (C.R. 244, 1460, 1957). Points of intensity 0.5 - 0.75 - 1.0 - 1.5 and 2.0 times 10^{-22} watts/m²/c/s are joined day after day in the form of isophotes. Black dots give the position of the center of the radio spots for each day; a line indicates the width of the recorded lobe pattern when it can be measured with certainty. For each radio spot the smoothed intensity around noon is given in 10^{-22} watts/m²/c/s.


Note that the isophotes cannot be measured when a radio spot of large intensity is on the disk.

Spectrum Observations

Data on solar radio emission in the spectral range 25-580 Mc/s recorded at the Radio Astronomy Station of Harvard College Observatory, Fort Davis, Texas, are presented. The research program is supported by financial assistance from the Air Force Cambridge Research Center, through the offices of Sacramento Peak Observatory.

The receiving equipment consists of five separate sweep frequency receivers covering the bands 25-50, 50-100, 100-180, 170-320, 300-580 Mc/s. The 25-50 and 50-100 Mc/s receivers are each connected to broad band dipoles which are cross polarised and mounted over a reflecting screen. The other three receivers are attached to separate broad band feeds mounted coaxially at the primary focus of an 8.55 meter diameter paraboloid, the 170-320 Mc/s feed being cross polarised with the other two feeds. The effective collecting area of the antenna is 40 sq. meters at 100 Mc/s and 45 sq. meters at 500 Mc/s.

The four types of recognized spectral activity are idealized below:



The large scale examples of continuum, sometimes called Type IV, are listed as "Cont. IV" in the tables. Photographic examples of the bursts have been published by Maxwell, Swarup, and Thompson (Proc. IRE 46, 142, 1958), and Maxwell (Sky and Telescope 17, 388, 1958; 18, 544, and 556, 1959). A few remaining solar radio bursts are tabulated as unclassified.

The symbols used in the tables are:

- b = single burst
- g = small group (<10) of bursts
- G = large group (≥ 10) of bursts
- = Arrows indicate continuity of solar activity between two Greenwich days.

The minimum detectable level of solar activity is a function of frequency: approximately 5×10^{-22} watts meter⁻² (c/s)⁻¹ at 500 Mc/s. The equipment records signals over an intensity range of approximately 10,000:1. There are three classes of intensity given in the tables. For 100 Mc/s they are:

- 1 = Faint, 5 to 40×10^{-22} watts meter⁻² (c/s)⁻¹.
- 2 = Moderate, 30 to 200×10^{-22} .
- 3 = Strong, $>200 \times 10^{-22}$.

The times are Universal Time (U. T.). The accuracy is to the nearest half minute, except in the case of major outbursts which are specified to the nearest 0.1 minute.

Details of the frequency ranges of activity may be obtained on request to the Radio Astronomy Station, Ft. Davis, Texas.

Y GEOMAGNETIC ACTIVITY INDICES

C, Kp, Ap, and Selected Quiet and Disturbed Days -- The data in the table are: (1) preliminary international character figures, C; (2) geomagnetic planetary three-hour range indices, Kp; (3) daily "equivalent amplitude," Ap; (4) magnetically selected quiet and disturbed days.

This table is made available by the Committee on Characterization of Magnetic Disturbances of IAGA, IUGG. The Meteorological Office, De Bilt, Holland collects the data from magnetic observatories distributed throughout the world, and compiles C and selected days. The Chairman of the Committee computes the planetary and equivalent amplitude indices. The same data are also published quarterly in the Journal of Geophysical Research along with data on sudden commencements (sc) and solar flare effects (sfe).

The C-figure is the arithmetic mean of the subjective classification by all observatories of each day's magnetic activity on a scale of 0 (quiet) to 2 (storm).

Kp is the mean standardized K-index from 12 observatories between geomagnetic latitudes 47 and 63 degrees. The scale is 0 (very quiet) to 9 (extremely disturbed), expressed in thirds of a unit, e.g. 5- is $4 \frac{2}{3}$, 5o is $5 \frac{0}{3}$, and 5+ is $5 \frac{1}{3}$. This planetary index is designed to measure solar particle-radiation by its magnetic effects, specifically to meet the needs of research workers in the ionospheric field. A complete description of Kp has appeared in Bulletin 12b, "Geomagnetic Indices C and K, 1948" of the Association of Terrestrial Magnetism and Electricity (IATME), International Union of Geodesy and Geophysics.

Ap is a daily index of magnetic activity on a linear scale rather than on the quasi-logarithmic scale of the K-indices. It is the average of the eight values of an intermediate 3-hourly index "ap," defined as one-half the average gamma range of the most disturbed of the three force components, in the three-hour interval at standard stations; in practice, ap is computed from the Kp for the 3-hour interval. The extreme range of the scale of Ap is 0 to 400. The method is described in IATME Bulletin No. 12h (for 1953) p. viii f. Values of Ap (like Kp and Cp) have been published for the Polar Year 1932/33 and for the years 1937 onwards.

The magnetically quiet and disturbed days are selected in accordance with the general outline in Terr. Mag. (predecessor to J. Geophys. Res.) 48, pp 219-227, December 1943. The method in current use calls for ranking the days of a month by their geomagnetic activity as determined from the following three criteria with equal weight: (1) the sum of the eight Kp's; (2) the sum of the squares of the eight Kp's; and (3) the greatest Kp.

Chart of Kp by Solar Rotations -- The graph of Kp by solar rotations is furnished through the courtesy of Dr. J. Bartels, Geophysikalisches Institute, Gottingen.

VI RADIO PROPAGATION QUALITY INDICES

One can take as the definition of a radio propagation quality index: the measure of the efficiency of a medium-powered radio circuit operated under ideal conditions in all respects, except for the variable effect of the ionosphere on the propagation of the transmittal signal. The indices given here are derived from monitoring and circuit performance reports, and are the nearest practical approximation to the ideal index of propagation quality.

Quality indices are usually expressed on a scale that ranges from one to nine. Indices of four or less are generally taken to represent significant disturbance. (Note that for geomagnetic K-indices, disturbance is represented by higher numbers.) The adjectival equivalents of the integral quality indices are as follows:

1 = useless	4 = poor-to-fair	7 = good
2 = very poor	5 = fair	8 = very good
3 = poor	6 = fair-to-good	9 = excellent

CRPL forecasts are expressed on the same scale. The tables summarizing the outcome of forecasts include categories P-Perfect; S-Satisfactory; U-Unsatisfactory; F-Failure. The following conventions apply:

P - forecast quality equal to observed	U - forecast quality two or more grades different from observed when <u>both</u> forecast and observed were > 5 , or both < 5
S - forecast quality one grade different from observed	F - other times when forecast quality two or more grades different from observed

Full discussion of the reliability of forecasts requires consideration of many factors besides the over-simplified summary given.

The quality figures represent a consensus of experience with radio propagation conditions. Since they are based entirely on monitoring or traffic reports, the reasons for low quality are not necessarily known and may not be limited to ionospheric storminess. For instance, low quality may result from improper frequency usage for the path and time of day. Although, wherever it is reported, frequency usage is included in the rating of reports, it must often be an assumption that the reports refer to optimum working frequencies. It is more difficult to eliminate from the indices conditions of low quality for reasons such as multipath or interference. These considerations should be taken into account in interpreting research correlations between the Q-figures and solar, auroral, geomagnetic or similar indices.

North Atlantic Radio Path -- The CRPL quality figures, Qa, are compiled by the North Atlantic Radio Warning Service (NARWS), the CRPL forecasting center at Ft. Belvoir, Virginia, from radio traffic data for North Atlantic transmission paths closely approximating New York-to-London. These are reported to CRPL by the Canadian Defense Research Board, Canadian Broadcasting Corporation, and the following agencies of the U.S. Government:--Coast Guard, Navy, Army Signal Corps, U.S. Information Agency. Supplementing these data are CRPL monitoring, direction-finding observations and field-strength measurements of North Atlantic transmissions made at Belvoir.

The original reports are submitted on various scales and for various time intervals. The observations for each 6-hour interval are averaged on the original scale. These 6-hour indices are then adjusted to the 1 to 9 quality-figure scale by a conversion table prepared by comparing the distribution of these indices for at least four months, usually a year, with a master distribution determined from analysis of the reports originally made on the 1 to 9 quality-figure scale. A report whose distribution is the same as the master is thereby converted linearly to the Q-figure scale. The 6-hourly quality figure is the mean of the reports available for that period.

The 6-hourly quality figures are given in this table to the nearest one-third of a unit, e.g. 5.0 is 5 and 0/3; 5- is 4 and 2/3; 5+ is 5 and 1/3. Other data included are:

(a) Whole-day radio quality indices, which are weighted averages of the four 6-hourly indices, with half weight given to quality grades 5 and 6. This procedure tends to give whole-day indices suitable for comparison with whole-day advance forecasts which seek to designate the days of significant disturbance or unusually quiet conditions.

(b) Short-term forecasts, issued every six hours by the North Atlantic Radio Warning Service. These are issued one hour before 00^h, 06^h, 12^h, 18^h, UT and are applicable to the period 1 to 7 hours ahead.

(c) Advance forecasts (CRPL-J) are issued once a week and are applicable 1 to 7 days ahead. They are modified as necessary by the Special Disturbance Warning (CRPL-SDW) applicable 1 to 3 days ahead, which may be followed by a supplementary forecast (CRPL-Js) applicable to days remaining until next CRPL-J forecast. The forecast entitled "final" consists of the most recent of the above forms and is scored against the whole-day quality index.

(d) Half-day averages of the geomagnetic K indices measured by the Fredericksburg Magnetic Observatory of the U.S. Coast and Geodetic Survey.

A chart compares the short-term forecasts with Qa-figures. A second chart compares the outcome of the final advance forecasts with a type of "blind" forecast. For the latter, the frequency for each quality grade, as determined from the distribution of quality grades in the four most recent months of the current season, is partitioned among the grades observed in the current month in proportion to the frequencies observed in the current month.

Ranges of useful frequencies on the North Atlantic radio path are shown in a series of diagrams, one for each day. The shaded area indicates the range of frequencies for which transmissions of quality 5 or greater were observed. The blacker the diagram, the quieter the day has been; a narrow strip indicates either high LUHF, low MUF, or both. These diagrams are based on data reported to CRPL by the German Post Office through the Fernmeldetechnischen Zentralamt, Darmstadt, Germany, being observations every one and a half hours of selected transmitters located in the eastern portion of North America. Since January 6, 1958 the transmitters monitored are restricted to those located north of 39° latitude. The magnetic activity index, A_{pr} , from Fredericksburg, Va., is also given for each day.

Note: Beginning with data for September 1955, Qa has been determined from reports that are available within a few hours or at most within a few days, including for the first time, the CRPL observations. Therefore these are the indices by which the forecasters assess every day the conditions in the recent past. Over a period of several years, they have closely paralleled the former Qa indices which excluded CRPL observations and included three additional reports received after a considerable lag. Qa was first published to the nearest one-third of a unit at the same time.

North Pacific Radio Path -- The CRPL quality figures, Qp, are compiled by the North Pacific Radio Warning Service (NPRWS), the CRPL forecasting center at Anchorage, Alaska, from radio traffic data for moderately long transmission paths in the North Pacific equivalent to Seattle-to-Anchorage or Anchorage-to-Tokyo. These include reports to CRPL by the Army Command and Administrative Network, U. S. Air Force and Federal Aeronautical Administration. In addition, there are CRPL monitoring, direction finder observations and field strength measurements of suitable transmissions.

The original reports are on various scales and for various time intervals. The observations for each 8 hours or 24 hour period are averaged on the original scale. This average is compared with reports for the same period in the preceding two months and expressed as a deviation from the 3-month mean. The deviations are put on the 1 to 9 scale of quality which is assumed to have a standard deviation of 1.25 and a mean for the various periods as follows:

07-18 hours UT	5.33	00-24 hours UT	5.67
19-06	6.00		

The 12-hour and 24-hour indices Qp are determined separately. Each index is a weighted mean where the CRPL observations have unit weight and the others are weighted by the correlation coefficient with the CRPL observations.

The table, analagous to that for Qa, includes the 12-hourly quality figures; whole day quality figures; short-term forecasts issued by NPRWS twice daily at 06^h and 18^h UT, applicable to the stated 12-hour periods; advance forecasts issued weekly by NPRWS (CRPL-Jp report) modified as necessary by Special Disturbance Warnings (CRPL-SDW) and supplementary forecasts (CRPL-Jps); and half-day averages of geomagnetic K indices from Sitka.

The chart compares the outcome of the final advance forecasts, on the same basis as the similar chart for the North Atlantic Radio Path.

Note: Beginning with March 15, 1959 the short-term forecast schedule was changed from three times daily to twice daily. The North Pacific quality figures used for evaluation are now 12-hourly rather than 8-hourly.

VII ALERT PERIODS AND SPECIAL WORLD INTERVALS


This table gives the Advance Geophysical Alerts as initiated by the Western Hemisphere Regional Warning Center at Ft. Belvoir, Va., and also the Worldwide Geophysical Alerts and Special World Intervals as designated by the World Warning Agency, Ft. Belvoir, Va.

Advance Alerts are of four types, defined as follows:

- 1 - Solar Flare Alert -- this warning is issued whenever a solar flare of median importance 2 plus or greater has been reported. There will be only one alert issued per flare and only one a day at most.
- 2 - Magnetic Storm Alert -- this warning is issued whenever a significant magnetic storm, K figure 5 or greater at a middle latitude station has begun.
- 3 - Cosmic Ray Alert -- this warning is issued whenever a very outstanding change in cosmic ray flux has been observed -- increase or decrease.
- 4 - Aurora Alert -- this warning is issued whenever a magnetic storm in middle latitudes has reached K figure 7 intensity or whenever selected auroral stations report the presence of outstanding aurora.

Worldwide Alerts are of the same types as the Advance Alerts, except that the Solar Flare Alert and Cosmic Ray Decrease Alert are omitted. Alert announcements include the event and time of event upon which the alert is based, and, in the case of the Advance Alerts, the station reporting the event.


The World Alerts and Special World Intervals are issued by the World Warning Agency on decisions based on Advance Alerts, advice received from Regional Warning Centers and overall policy.



DAILY SOLAR INDICES

Dec. 1959	American Relative Sunspot Numbers R_A'
1	164
2	192
3	194
4	174
5	155
6	152
7	131
8	138
9	97
10	76
11	72
12	74
13	79
14	82
15	120
16	108
17	105
18	131
19	163
20	174
21	154
22	116
23	110
24	121
25	143
26	134
27	150
28	153
29	120
30	113
31	131
Mean:	129.9

Jan. 1960	Zürich Provisional Relative Sunspot Numbers R_Z	Daily Values Solar Flux at 2800 Mc, Ottawa, Canada Flux
1	136	
2	110	175
3	133	182
4	156	193
5	158	213
6	174	215
7	167	224
8	153	219
9	150	201
10	127	194
11	143	200
12	108	184
13	108	178
14	118	176
15	112	183
16	119	183
17	117	179
18	89	176
19	80	164
20	94	157
21	103	162
22	134	172
23	138	188
24	130	210
25	152	230
26	209	242
27	186	248
28	159	252
29	193	237
30	178	230
31	178	224
Mean:	139.1	199.7



CALCIUM PLAGGE AND SUNSPOT REGIONS

JANUARY 1960

CMP Jan. 1960	Lat	McMath Plage Number	Return of Region	Calcium Plage Data				Sunspot Data		
				CMP Values		History, Age		CMP Values		History
				Area	Int.			Area	Count	
01.1	N09	5512	5478	2000	2.5	<i>l - l</i>	2	80	7	<i>b / l</i>
02.1	N22	5513	5478	3000	2.5	<i>l \ l</i>	2	50	2	<i>l \ d</i>
03.2	N08	5519	New	400	1	<i>b ^ d</i>	1			
04.3	S15	5514	5482	3200	2.5	<i>l - d</i>	5	80	4	<i>l - l</i>
04.5	N11	5516	New	1400	2.5	<i>l / l</i>	1	20	1	<i>l \ l</i>
05.6	N09	5517	5484	700	2.5	<i>l - l</i>	5	320	6	<i>l - l</i>
06.6	S18	5515	5486	3500	3	<i>l - l</i>	4	370	3	<i>l - l</i>
07.6	N07	5521	5495	900	2	<i>l - l</i>	2	50	2	<i>b ^ d</i>
07.7	N25	5520	New	2700	2.5	<i>l - l</i>	1	20	1	<i>l / l</i>
09.0	S23	5522	New	3200	3	<i>l - l</i>	1	440	2	<i>l - l</i>
09.1	N17	5524	New	600	2.5	<i>l - l</i>	1	(20)	(1)	<i>l \ d</i>
09.3	S01	5523	New	2100	3	<i>l - l</i>	1	560	14	<i>l - l</i>
10.2	S16	5525	New	6300	3	<i>l - l</i>	1	1020	6	<i>l \ l</i>
10.2	S03	5526	5490	600	2.5	<i>l / l</i>	4	70	3	<i>b ^ d</i>
11.6	N20	5527	5491	3500	2.5	<i>l - l</i>	2	730	5	<i>l - l</i>
12.5	S21	5530	New	500	1	<i>l / l</i>	1	50	1	<i>b / l</i>
13.4	N03	5528	*	3000	2	<i>l \ l</i>	6,2			
14.4	N25	5531	New	500	1.5	<i>l \ d</i>	1			
15.1	S16	5529	New	300	1.5	<i>l \ d</i>	1			
15.9	N10	5532	5497	600	2	<i>l \ d</i>	6			
16.3	S06	5533	5499	800	2.5	<i>l / l</i>	3	50	1	<i>l \ d</i>
16.7	N22	5535	New	500	2	<i>l \ d</i>	1			
17.7	N09	5534	5501	2000	2.5	<i>l - l</i>	4			
17.8	S18	5536	5500	800	1	<i>l / l</i>	4	60	1	<i>l \ d</i>
19.3	N07	5538	New	2200	3.5	<i>l - l</i>	1	100	5	<i>l - l</i>
19.5	N20	5537	5502	1700	2	<i>l - l</i>	4			
21.1	N09	5540	**	3700	3	<i>l - l</i>	3	150	4	<i>l \ d</i>
21.5	N26	5539	**	2600	3	<i>l - l</i>	3	540	4	<i>l - l</i>
22.2	N13	5541	**	2900	3	<i>l - l</i>	3	(20)	(1)	<i>l \ d</i>
23.0	S13	5543	New	700	2	<i>l - l</i>	1	100	3	<i>b ^ d</i>
23.3	N26	5542	**	3200	3	<i>l - l</i>	3			
23.4	N11	5545	**	2900	3	<i>l - l</i>	3	390	1	<i>l - l</i>
25.0	N12	5546	5507	(5600)	(2.5)	<i>l - l</i>	10			
26.5	S13	5547	5510	(2000)	(2)	<i>l - l</i>	2	190	1	<i>l - l</i>
26.6	N24	5548	5509	4900	3	<i>l - l</i>	4	610	5	<i>l - l</i>
27.2	N09	5549	***	2000	3	<i>l - l</i>	3	400	10	<i>l - l</i>
29.5	N12	5550	***	11000	3	<i>l - l</i>	3	580	26	<i>l - l</i>
31.8	S18	5551	5514	5000	3	<i>l - l</i>	6	510	12	<i>l - l</i>

* 5493 and 5494.
 ** 5504, 5505, 5506.
 *** 5511, 5512, 5513.

COMMERCE - STANDARDS - BOULDER

CORONAL LINE EMISSION INDICES

JANUARY 1960

CMP Jan 1960	North East Quadrant (observed 7 days earlier)			South East Quadrant (observed 7 days earlier)			South West Quadrant (observed 7 days later)			North West Quadrant (observed 7 days later)				
	G ₆	G ₁	R ₁	G ₆	G ₁	R ₁	G ₆	G ₁	R ₁	G ₆	G ₁	R ₁		
1	x	x	x	x	x	x	75	124	19	48	161	199	30	54
2	x	x	x	x	x	x	56	85	7	8	92	109	13	16
3	x	x	x	x	x	x	x	x	x	x	x	x	x	x
4	93	120	20	91	115	57	x	x	x	x	x	x	x	x
5	141	167	15	148	370	51	x	x	x	x	x	x	x	x
6	162	194	29	146	190	75	x	x	x	x	x	x	x	x
7	x	x	x	x	x	x	x	x	x	x	x	x	x	x
8	x	x	x	x	x	x	127	182	41	65	131	167	32	41
9	x	x	x	x	x	x	x	x	x	x	x	x	x	x
10	88	102	x	99*	126	x	x	x	x	x	x	x	x	x
11	117	132	24	x	x	24	x	x	x	x	x	x	x	x
12	x	x	x	x	x	x	87	135	22	36	113	137	41	69
13	x	x	x	x	x	x	x	x	x	x	x	x	x	x
14	100	114	x	55	78	x	59	99	23	39	84	107	25	39
15	110	128	25	74	104	20	76	98	12	20	82	99	14	24
16	78	107	17	57	78	10	x	x	x	x	x	x	x	x
17	x	x	x	x	x	x	52	114	x	x	107	174	x	x
18	x	x	x	x	x	x	x	x	x	x	x	x	x	x
19	x	x	x	x	x	x	30	47	13	16	137	208	47	63
20	x	x	x	x	x	x	x	x	x	x	x	x	x	x
21	x	x	x	x	x	x	x	x	x	x	x	x	x	x
22	124*	156	52	47	54	15	x	x	x	x	x	x	x	x
23	x	x	x	x	x	x	x	x	x	x	x	x	x	x
24	x	x	x	x	x	x	x	x	x	x	x	x	x	x
25	x	x	x	x	x	x	x	x	x	x	x	x	x	x
26	123	138	21	93	135	24	x	x	x	x	x	x	x	x
27	x	x	x	x	x	x	x	x	x	x	x	x	x	x
28	132*	152	40	61	104	39	x	x	x	x	x	x	x	x
29	143	178	30	51	117	20	x	x	x	x	x	x	x	x
30	x	x	x	x	x	x	x	x	x	x	x	x	x	x
31	219	267	x	157	261	x	x	x	x	x	x	x	x	x

COMMERCE - STANDARDS - BOULDER

x = no observations. a = index computed from low weight data. * = yellow line observed.

SOLAR FLARES

JANUARY 1960

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME			LOCATION			DURA- TION — MINUTES	IM- POR- TANCE	OBS. COND.	MEASUREMENTS				PROVISIONAL IONOSPHERIC EFFECT
		START	END	MAX. PHASE	APPROX. LAT.	MAGN. BLAZE REGION	TIME — U T				MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	MAX. WIDTH Rg	MAX. INT. %	
HAWAII	01 0128	0200 D	0158		N12 W54	5507	32 D	1	3	0158	1.40				
MCMATH	01 1607	1650			N24 E80	5520	43	1	1	1618					
WENDEL	02 0950	E	1025 D		N17 W20	5511	35 D	1			3.00				
WENDEL	02 1029	1040 D			N13 E64	5520	11 D	1			3.00				
WENDEL	02 1320	1333 D			S21 E55	5515	13 D	1			3.00				
HAWAII	03 1944	E	2010		N17 W28	5513	26 D	16	3	1947	2.10				
HAWAII	03 1948	1952		1948	S23 E85	5525	4	1	3	1948	1.10				
CAPRI S	04 0327	0906			S19 E77	5525	39	26	1	0852	1.90	6.70			
ARCETRI	04 0847	0854			S14 E77	5525	7	16	3	0848	1.65	4.95			
HAWAII	04 1956	E	2014		N08 W38	5512	18 D	1	2	2004	1.10				
{ LOCKHEED	04 2025	2058		2029	N08 W37	5512	33	1	1	2029	1.90				30
{ HAWAII	04 2026	2048		2030	N09 W37	5512	22	1	3	2030	1.10				
{ LOCKHEED	04 2142	2207		2149	N15 W57	5511	25	1	1	2149	2.20				10
ARCETRI	05 0920	E	0959 D		N10 E04	5517		1	3	0920	2.50				
{ ARCETRI	05 0955	1007			S11 E65	5525	4 D	1	3			3.00			
{ WENDEL	05 0955	1007			S12 E58	5525	12	1							
ARCETRI	07 0809	E	0819 D		S12 E40	5525	10 D	1	3						
ARCETRI	07 0819	E	1542 D		N08 W71	5512		1	3						
CAPRI S	07 1504	E			N07 W78	5512	38 D	3	3	1514	3.70	13.00			S-SWF
{ LOCKHEED	08 1750	1950		1810	S23 W33	5515	120	1	2	1825	2.50				20
{ LOCKHEED	08 1750	1950		1825	S23 W33	5515	120	1	2	1825	2.50				20
{ LOCKHEED	08 1800	2050		1840	N28 W12	5520	170	2	2	1840	6.40				30
{ SAC PEAK	08 1806	2050 U		1842	N26 W14	5520	164 U	2	3		6.85				19
{ HAWAII	08 1820	E	2048		N28 W14	5520	148 D	2	2	1850	3.10				
SAC PEAK	09 2106	2130		2112	N19 E21	5527	24	1	2			2.76			13
ARCETRI	10 0811	0815			S17 W50	5515	4	1	3						
ARCETRI	10 0907	E	0925 D		N08 W90	5516	18 D	1	3						
WENDEL	10 1139	E	1210 D		N08 W85	5516	31 D	16				5.00			
{ LOCKHEED	11 2040	U	2355 D		N23 E03	5527	195 D	3	1	2128	22.00				30
{ HAWAII	11 2058	E	2326		N22 E02	5527	148 D	3	3	2124	10.30				Slow S-SWF
HAWAII	11 2244	2422		2250	N18 W08	5527	98	16	3	2250	1.90				
HAWAII	11 2258	2312		2302	S12 W90	5515	14	1	3	2302	.50				
SAC PEAK	12 1646	1710		1650	S10 W37	5525	24	1	3			2.60			17
CAPRI S	13 1216	E	1232 D		S18 W44	5525	16 D	1	1	1217	1.70				
LOCKHEED	13 2247	2350		2305	S03 W08	5530	63	1	1	2305	2.10				10
{ WENDEL	15 1334	1400 D			S20 W66	5525	26 D	2	2	1401	4.80				Slow S-SWF
{ CAPRI S	15 1336	1455 D			S20 W71	5525	79 D	2	2	1404	11.50				2.50
ONDREJOV	15 1400	E	1412 D		S20 W69	5525	12 D	2	2	1658	2.00				20
LOCKHEED	15 1645	1730 U		1658	S06 E08	5553	45	1	1	1658	2.00				30
LOCKHEED	15 1730	D	1738 U		N09 E90	5540	8 D	1	1	1732	2.00				Slow S-SWF
WENDEL	16 1005	1021 D			N26 E73	5539	16 D	1	1			3.00			

SOLAR FLARES

JANUARY 1960

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME			LOCATION			DURA- TION — MINUTES	IM- POR- TANCE	ORS. COND.	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT
		START	END	MAX. PHASE	APPROX. LAT.	MER. DIST.	MONTH PLAGE REGION				TIME U - T	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	
{ HAWAII LOCKHEED	16	2108	2112 D	2112	N26 E62	5539	4 D	1	2		1.40			Slow S-SWF
	16	2239	2335	2252	N11 E76	5540	56	2	2		5.00			
	16	2240	2258 D	2248	N13 E76	5540	18 D	1	2		1.50			
{ HAWAII LOCKHEED	17	0032	0046 D	0036	N11 E67	5545	14 D	1	2		1.10			Slow S-SWF
	17	1708 E	1900 U	1720	N17 E62	5545	112 D	16	1		4.20			
	17	1708 E	1900 U	1750	N17 E62	5545	112 D	16	1		4.20			
	17	1800 E	1856		N23 E72	5542	56 D	16	2		2.20			
ARCETRI	18	0934 E	0943 D		N08 E41	5541	9 D	1	3		1.00			G-SWF
	18	2250	2358	2308	N17 E58	5545	68	1	3		1.40			
	18	2314	2442		N16 E49	5541	88	1	3		1.40			
{ SAC PEAK LOCKHEED	19	1928	2028	1944	N14 E46	5545	60	1	2		3.97			G-SWF
	19	1947 E	2115 U	1950 U	N18 E45	5545	88 D	1	1		5.00			
WENDEL HAWAII	20	1038 E	1146 D		N07 W57	5528	68 D	1	3		1.30	3.00		
	20	2300	2316	2302	N12 E25	5541	16	1	3		1.90			
{ HAWAII ONDREJOV	23	0000	0014	0002	N27 W16	5539	14	16	3		0.002			2.90 2.70 4.80
	23	1040 E	1044 D		N09 W59	5538	4 D	1	3		1.041			
	23	1056 E	1104		N09 W59	5538	8 D	1	3		1058			
{ WENDEL ONDREJOV	23	1117	1130	1122	N09 W60	5538	13	1	3		1.122			17 30
	23	2038	2046	2038	N12 E49	5549	8	1	2		1.20			
{ ONDREJOV ONDREJOV	24	0855	0904	0858	N06 E57	5550	9	1	3		0.858			3.20 3.20 2.30
	24	0928 E	0941		N08 W69	5538	13 D	1	3		0.938			
	24	1241	1308 D		N06 E55	5550	27 D	1	3		1.245			
{ WENDEL ONDREJOV	24	1242 E	1300 D		N08 E56	5550	18 D	16	2			6.00		14.00 4.80
	24	1306	1358 D		N09 E38	5549	52 D	2	3		1.313			
	24	1309 E	1335 D		N08 E35	5549	26 D	2	2		4.00			
CAPRI S	24	1312 E	1402 D		N08 E35	5549	50 D	16	2			4.80		2.30
	25	1241	1303	1243	N06 E55	5550	22	1	3		1.40			
HAWAII	25	2004	2032	2008	N12 E50	5550	28	1	2					16
	26	1022 E	1028 D		N05 E31	5550	6 D	1	2					
{ SAC PEAK CAPRI S	27	1518	1544	1528	N01 E12	5549	26	1	2		2.60			16
	27	1520 E	1544 D		N05 E13	5549	24 D	1	2		3.00	3.10		
CAPRI S	28	0828 E	0954 D		N07 E12	5550	86 D	1	1		5.00			16
	28	2036	2118	2044	N00 W05	5550	42	16	2		4.78	5.00		
CAPRI S	29	1038 E	1045 D		S15 E27	5551	7 D	1	1		2.00			20 16 30
	29	1738	1810	1746	N06 W17	5550	32	1	2		2.40			
	29	1820	1902	1846	S19 E58	5554	42	1	2		2.43			
LOCKHEED HAWAII	30	0005	0033	0015	N03 W22	5550	28	1	2		2.00			30
	30	0110 E	0118		N10 W14	5550	8 D	1	2		1.40			
	30	0955 E	0922 D		S15 E21	5551	27 D	1	2					
{ WENDEL ARCETRI	30	0903 E	0926 D		S17 E17	5551	23 D	1	2			3.00		3.00
	30	0927 E	0932 D		S18 E49	5554	5 D	1	2					

SOLAR FLARES

JANUARY 1960

OBSERVATORY	DATE JAN 1960	OBSERVED UNIVERSAL TIME		LOCATION		DURA- TION — MINUTES	IM- POR- TANCE	OBS. COND.	TIME — U T	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT	
		START	END	APPROX. LAT.	APPROX. MER. DIST.					MEMBER PLAGE REGION	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.		MAX. WIDTH Ec
WENDEL	30	1035	1057	N09 E36	5552	22	1			3.00	4.00			
	30	1045	1101	S15 E20	5551	16	D			4.00				
	30	1138	1150	N09 W21	5550	12	D			3.00				
	30	1140	1152	N16 E19	5552	12	D			4.00				
	30	1139	1229	S16 E20	5551	30	D			4.00				
	{ CAPRI S	30	1355	1442	S17 E19	5551	163	D	3	1420	2.00	8.00		
		30	1306	1524	S17 E21	5551	89	D	3	1310	3.00	2.20		
	{ WENDEL	30	1306	1320	N03 W27	5550	14	D			5.00	3.30		
		30	1334	1321	N03 W26	5550	15	D			5.00	5.00		
	CAPRI S	30	1432	1502	N12 E35	5552	30	D	3	1436	3.00	3.70		
CAPRI S	31	1028	1042	N18 W17	5550	14	D	1	1031	2.00	2.10			
CAPRI S	31	1429	1509	S16 E04	5551	40	D	3	1451	4.80	5.00			
CAPRI S	31	1539	1605	S16 E04	5551	26	D	1	1600	3.00	3.10			
LOCKHEED	31	1913	2000	S18 E02	5551	47	D	1	1924	2.50			20	
LOCKHEED	31	2035	2103	S18 W00	5551	28	D	1	2047	3.20			20	


CAPRI G ANACAPRI - GERMAN MOSCOW-G MOSCOW - GAISH
 CAPRI S ANACAPRI - SWEDISH R O EDIN ROYAL OBSERVATORY, EDINBURGH
 GOOD HOPE ROYAL OBSERVATORY, CAPE OF GOOD HOPE R O HERST GREENWICH ROYAL OBSERVATORY, HERSTMONCEUX
 KIEV* KIEV UNIVERSITY SAC PEAK SACRAMENTO PEAK
 KODAIKANAL SCHAIJINS SCHAIJINSLAND
 KRASNAYA KRASNAYA PAKHRA USNRL UNITED STATES NAVAL RESEARCH LABORATORY
 LOCKHEED LOS ANGELES

SAC PEAK: ALL VALUES IN MAX. INT. COLUMN ARE ARBITRARY UNITS (0-40), NOT PERCENT OF CONTINUOUS SPECTRUM.
 E - LESS THAN & - PLUS
 D - GREATER THAN - - MINUS
 U - APPROXIMATE □ - NOT REPORTED

LOCKHEED OBSERVATIONS: ALL VALUES IN THE MAXIMUM INTENSITY COLUMN ARE ARBITRARY UNITS ON A SCALE OF 10 TO 40 - NOT PERCENT OF THE CONTINUOUS SPECTRUM.

INTERVALS OF NO FLARE PATROL OBSERVATIONS
 JANUARY 1960

III d



Stations Include:

- | | | |
|--------------------|----------|-----------------|
| Anacapri (Swedish) | Hawaii | Meudon |
| Arcetri | Lockheed | Ondrejov |
| Dunsink | McMath | Sacramento Peak |

COMMERCE - STANDARDS - BOULDER 820441-105-18

Noted as follows: Date-Universal Time-Coordinates

DECEMBER 1959

SAC PEAK	01	1504	N15 W75	* HAWAII	05	1946 E	N11 W04	SAC PEAK	13	1654	N15 E20
LOCKHEED	01	1649	N20 E16	LOCKHEED	05	2002	N07 E50	LOCKHEED	14	2017	S14 E11
LOCKHEED	01	1806	N24 W85	HAWAII	05	2008	N13 W10	SAC PEAK	14	2018	S13 E10
SAC PEAK	01	1808	N25 W90	LOCKHEED	05	2014	N10 W06	LOCKHEED	14	2034	S14 E11
SAC PEAK	01	1808	N11 E52	HAWAII	05	2026	N10 E01	LOCKHEED	14	2034	S14 E11
LOCKHEED	01	1808	N10 E54	LOCKHEED	05	2028	N09 E01	LOCKHEED	14	2034	S14 E11
LOCKHEED	01	1808	N10 E54	SAC PEAK	05	2038	N12 W12	LOCKHEED	14	2034	S15 W77
LOCKHEED	01	2008	N11 E51	LOCKHEED	05	2038	N10 W12	LOCKHEED	14	2304	S15 W77
LOCKHEED	01	2015	N22 E16	HAWAII	05	2040	N12 W12	HAWAII	15	0056 E	S08 E33
SAC PEAK	01	2018	N24 E18	LOCKHEED	05	2058	N09 E00	WENDEL	15	0823 E	S03 E23
LOCKHEED	01	2036	N10 E52	LOCKHEED	05	2058	N09 E02	* ARCTERI	15	0847 E	N11 E59
HAWAII	01	2038 E	N08 W07	LOCKHEED	05	2118	N06 W12	* WENDEL	15	1018 E	N23 W00
LOCKHEED	01	2041	N12 E02	SAC PEAK	05	2122	N07 W11	WENDEL	15	1119 E	N16 E02
SAC PEAK	01	2046	N10 W03	LOCKHEED	05	2137	N11 W05	* WENDEL	15	1120 E	N18 W04
SAC PEAK	01	2126	N09 W18	SAC PEAK	05	2138	N11 W04	WENDEL	15	1129 E	S03 E22
SAC PEAK	01	2150	N08 W08	LOCKHEED	05	2154	N10 W05	WENDEL	15	1201 E	N09 E56
LOCKHEED	01	2156	N09 W11	LOCKHEED	05	2154	N10 W05	LOCARNO	15	1320	N20 W04
HAWAII	01	2214	N13 W12	HAWAII	05	2200 E	N11 W12	WENDEL	15	1340 E	S15 E50
LOCKHEED	01	2228	N24 W85	LOCKHEED	05	2220	N10 W15	LOCKHEED	15	1634	N09 W90
LOCKHEED	01	2252	N09 W09	LOCKHEED	05	2223	N11 W06	LOCKHEED	15	1652	S06 E31
HAWAII	01	2306 E	N14 W01	LOCKHEED	05	2318	N10 W15	LOCKHEED	15	1656	N08 E54
* LOCKHEED	01	2333	N09 W12	HAWAII	05	2320 E	N13 W14	LOCKHEED	15	1713	N09 W90
HAWAII	02	0008 E	S03 E53	* HAWAII	05	2342	N09 W09	LOCKHEED	15	1800	N16 W06
WENDEL	02	1341 E	N11 E38	LOCARNO	06	0955	N11 W12	LOCKHEED	15	1826	N06 W56
SAC PEAK	02	1530	N11 E36	LOCARNO	06	1200	N09 W20	LOCKHEED	15	1835	N16 W07
SAC PEAK	02	1554	S10 W18	LOCARNO	06	1226	N11 W13	LOCKHEED	15	1927	S18 E75
LOCKHEED	02	1554	N07 W14	* WENDEL	06	1233 E	N10 W08	LOCKHEED	15	1948	N25 W07
LOCKHEED	02	1619	N10 E39	LOCARNO	06	1238	N08 W18	LOCKHEED	15	1955	S08 E29
LOCKHEED	02	1619	N10 E39	LOCARNO	06	1254	N11 W14	LOCKHEED	15	1956	S17 E85
SAC PEAK	02	1622 D	N10 E38	LOCARNO	06	1350	N07 W63	LOCKHEED	15	2007	S08 E29
LOCKHEED	02	1626	N09 W17	SAC PEAK	06	1358	N07 W63	HAWAII	15	2154	N17 W08
SAC PEAK	02	1638	N09 W18	SAC PEAK	06	1526	N07 W23	LOCKHEED	15	2154	N16 W10
LOCKHEED	02	1639	N11 E41	SAC PEAK	06	1538	N08 W19	LOCKHEED	15	2358	S03 E18
SAC PEAK	02	1646	N11 E41	SAC PEAK	06	1556	N13 W13	* ARCTERI	16	0846 E	N05 W66
WENDEL	02	1648	S15 E74	SAC PEAK	06	1611	N07 W22	* WENDEL	16	0913 E	N05 W66
LOCKHEED	02	1658	N08 W14	LOCKHEED	06	1618	N07 W22	WENDEL	16	0953 E	N03 W60
LOCKHEED	02	1711	S15 E74	SAC PEAK	06	1634	N09 W11	MCNATH	16	1509	S05 E55
SAC PEAK	02	1712	S15 E73	LOCKHEED	06	1635	N09 W11	LOCKHEED	16	1647	S15 W16
LOCKHEED	02	1748	N10 W26	LOCKHEED	06	1706	N09 W11	SAC PEAK	16	1722	S08 W32
SAC PEAK	02	1754	N10 E34	SAC PEAK	06	1710	N08 W77	LOCKHEED	16	1722	S07 W33
LOCKHEED	02	1754	N11 E33	* LOCKHEED	06	1714	N08 W20	LOCKHEED	16	1751	S16 E85
HAWAII	02	1756 E	N01 E35	LOCKHEED	06	1715	N09 W11	LOCKHEED	16	1819	S08 W33
* LOCKHEED	02	1758	N07 W20	SAC PEAK	06	1742	N09 W11	LOCKHEED	16	1838	S18 E66
* SAC PEAK	02	1800	N08 W19	LOCKHEED	06	1812	N11 W25	MCNATH	16	1840 E	S16 E85
SAC PEAK	02	1814	N12 E41	LOCKHEED	06	1812	N11 W25	HAWAII	16	1846 E	S21 E65
LOCKHEED	02	1815 E	N11 E41	SAC PEAK	06	1822	N34 E04	LOCKHEED	16	2300 E	S21 E65
* LOCKHEED	02	1815 E	N09 W12	SAC PEAK	06	1822	N36 E03	* WENDEL	17	0917 E	N11 E30
HAWAII	02	1826 E	N04 E39	* LOCKHEED	06	1903	N11 W13	WENDEL	17	1018 E	N11 E30
* SAC PEAK	02	1844	N11 W14	SAC PEAK	06	1938	N08 W21	LOCARNO	17	1120	S16 E53
* LOCKHEED	02	1850 E	N10 W13	LOCKHEED	06	1945	S19 W90	MCNATH	17	1547	N15 W36
SAC PEAK	02	2108	N09 W21	SAC PEAK	06	1954	N11 W26	MCNATH	17	1615 E	S04 W07
LOCKHEED	03	0011	N10 W16	LOCKHEED	06	1954	N11 W26	LOCKHEED	17	1724	S03 W47
WENDEL	03	0950 E	N19 E7	SAC PEAK	06	2100	N10 W22	LOCKHEED	17	1739	S06 W47
ARCTERI	03	0957 E	N08 W29	LOCKHEED	06	2100	N09 W23	LOCKHEED	17	1801	S19 E50
WENDEL	03	1012 E	N12 E26	LOCKHEED	06	2140	N13 W25	LOCKHEED	17	1816	S07 W47
CAPRI S	03	1237 E	N08 W25	SAC PEAK	06	2140	N12 W24	LOCKHEED	17	1824	S04 W07
* CAPRI S	03	1312	N08 W25	LOCKHEED	06	2152	N09 W14	SAC PEAK	17	1828	N08 W08
MCNATH	03	1516 E	N10 W20	SAC PEAK	06	2223	N12 W61	LOCKHEED	17	1939	N22 E73
MCNATH	03	1535 E	N31 E46	LOCKHEED	06	2302	N11 W27	* HAWAII	17	2144	S05 W46
* LOCKHEED	03	1607	N09 W33	LOCKHEED	06	2302	N11 W27	* SAC PEAK	17	2144	S06 W47
LOCKHEED	03	1615 E	S14 E64	LOCKHEED	07	0007	N07 W26	SAC PEAK	17	2209	S17 E52
MCNATH	03	1630 E	N10 E26	WENDEL	07	1243 E	N10 W33	CAPRI S	18	1343	N02 W15
LOCKHEED	03	1632	N09 E27	WENDEL	07	1314 E	N09 W19	SAC PEAK	18	1538	N23 E61
LOCKHEED	03	1632	N09 E27	SAC PEAK	07	1510	N10 W23	SAC PEAK	18	1546	S18 E40
LOCKHEED	03	1634	N13 W36	SAC PEAK	07	1610	N10 W24	MCNATH	18	1548	S18 E40
LOCKHEED	03	1746	N13 W36	HAWAII	07	1820 E	N16 W57	LOCKHEED	18	1548	S18 E40
LOCKHEED	03	1749	N10 E27	HAWAII	07	2036	E13	LOCKHEED	18	1548	S18 E40
HAWAII	03	1902	N07 E23	HAWAII	07	2212	N25 E50	LOCKHEED	18	1550 E	N21 W45
LOCKHEED	03	1903	N10 E24	WENDEL	08	1154 E	N29 E36	LOCKHEED	18	1814 E	N26 E63
LOCKHEED	03	1910	S14 E62	WENDEL	08	1425	N13 W45	LOCKHEED	18	1816	N34 W24
LOCKHEED	03	1944	N09 E24	MCNATH	08	1440 E	N17 W01	SAC PEAK	18	1816	N30 W20
LOCKHEED	03	2006	N09 W39	MCNATH	08	1441	N28 E36	SAC PEAK	18	1926	N10 E13
LOCKHEED	03	2011	N08 W35	MCNATH	08	1444	S17 W07	HAWAII	18	1940 E	N10 E15
LOCKHEED	03	2018	N09 E22	MCNATH	08	1530	N07 W49	LOCKHEED	18	2121	N06 W64
LOCKHEED	03	2042	N08 W39	MCNATH	08	1621	N13 W43	HAWAII	18	2254	N20 E60
LOCKHEED	03	2103	N12 E25	MCNATH	08	1656 E	N28 E34	HAWAII	19	0014	S05 W64
LOCKHEED	03	2142	N08 W35	WENDEL	09	1245 E	N11 W54	HAWAII	19	0030	N02 W27
LOCKHEED	03	2201	S16 E62	* CAPRI S	09	1317	N10 W48	HAWAII	19	1425	N23 E48
LOCKHEED	03	2228	N10 E22	* SAC PEAK	09	1626 E	N10 W55	MCNATH	19	1548	N24 E48
LOCKHEED	03	2250 U	N08 W37	WENDEL	10	1222 E	N09 W15	MCNATH	19	1756 E	N25 E48
HAWAII	04	0146	N02 E20	MCNATH	10	1455	N18 E62	WENDEL	19	1805 E	N11 W03
SAC PEAK	04	1516	N12 W43	LOCKHEED	10	1548	N08 W74	HAWAII	19	1916	S01 W24
LOCKHEED	04	1611	N13 W43	LOCKHEED	10	1559	N20 W27	HAWAII	19	1918	S16 W43
LOCKHEED	04	1621	N10 E16	LOCKHEED	10	1610	N09 W74	HAWAII	19	2114	N11 W01
SAC PEAK	04	1624	N08 E13	LOCKHEED	10	1706	N21 W28	* SAC PEAK	19	2202 E	N25 E45
LOCKHEED	04	1624	N07 E13	LOCKHEED	10	1718	N10 W72	HAWAII	19	2306	N11 W01
LOCKHEED	04	1641	N08 W48	LOCKHEED	10	1938	S06 E47	HAWAII	19	2320 E	N08 E31
LOCKHEED	04	1701	N07 E19	* HAWAII	10	2034	N13 E68	* HAWAII	20	0010	N25 E43
LOCKHEED	04	1741	N06 W49	HAWAII	10	2132 E	N14 W77	* CAPRI S	20	0912	S17 E07
LOCKHEED	04	1822	N09 E14	LOCKHEED	10	2344	N13 W84	SAC PEAK	20	2002	N16 W78
LOCKHEED	04	1840	S16 E51	LOCKHEED	10	2345	N07 W90	SAC PEAK	20	2022	N20 E38
LOCKHEED	04	1843	N08 E11	WENDEL	11	1112 E	N22 E53	SAC PEAK	20	2030	S04 W52
SAC PEAK	04	1846	N08 E11	WENDEL	11	1216 E	N09 W76	HAWAII	21	0142 E	N17 W80
SAC PEAK	04	1902	N10 E03	LOCKHEED	11	1625	N21 E48	* HAWAII	21	0145 E	N21 E32
LOCKHEED	04	1905	N11 E03	LOCKHEED	11	1651	N17 E49	LOCKHEED	21	1944	N12 E85
LOCKHEED	04	1915	N13 E10	LOCKHEED	11	1722	N11 W87	SAC PEAK	21	2054	N17 E08
LOCKHEED	04	1937	N10 E02	LOCKHEED	11	1754	S02 E77	LOCKHEED	21	2056	N17 E07
* LOCKHEED	04	1953	N08 W49	LOCKHEED	11	1920	S03 E71	LOCKHEED	21	2059	N25 E26
LOCKHEED	04	2008	N11 E02	SAC PEAK	11	2032	N16 E40	LOCKHEED	21	2100	N11 W22
LOCKHEED	04	2018	N11 E10	LOCKHEED	11	2039	N05 W85	LOCKHEED	21	2106	S18 W05
* LOCKHEED	04	2023	N08 W46	* LOCKHEED	11	2118	S02 E69	LOCKHEED	21	2126	N24 E23
SAC PEAK	04	2028	N11 E02	LOCKHEED	11	2145	N22 E47	SAC PEAK	21	2128	N20 E24
LOCKHEED	04	2044	N11 E02	HAWAII	11	2220 E	N12 W90	LOCKHEED	21	2142	N11 E85
LOCKHEED	04	2114	N10 E02	LOCKHEED	11	2238	N11 W38	HAWAII	21	2336 E	N12 W01
LOCKHEED	04	2127	N08 E09	HAWAII	11	2256	N13 W38	LOCARNO	22	1110	N07 W08
LOCKHEED	04	2136	N11 E01	ARCTERI	12	0830 E	S20 W12	MCNATH	22	1502 E	N10 W07
LOCKHEED	04	2152	N09 E08	* ARCTERI	12	0831 E	N20 E40	MCNATH	22	1611	N10 W10
LOCKHEED	04	2233	N11 E00	WENDEL	12	0850 E	N15 E40	SAC PEAK	22	1612	N11 W09
LOCKHEED	04	2235	N08 E07	WENDEL	12	0950 E	N07 W43	SAC PEAK	22	1800	N13 W08
LOCKHEED	04	2318	N11 E00	SAC PEAK	12	1634	N17 E33	MCNATH	22	1804	N11 E15
LOCKHEED	04	2327	N09 E05	LOCKHEED	12	1713	N18 E35	SAC PEAK	22	1808	N12 W47
LOCARNO	05	1300 E	N10 W10	LOCKHEED	12	1906	N18 E32	SAC PEAK	22	1848	N15 W13
LOCARNO	05	1335 E	N09 W01	LOCKHEED	12	1917	N10 E98	MCNATH	22	1849	N12 W13
LOCARNO	05	14									

SUBFLARES

III f

Noted as follows: Date-Universal Time - Coordinates
DECEMBER 1959

* HAWAII	26	1826 E	N28 W47	SAC PEAK	28	1752	N17 W18	CAPRI S	30	0824 E	S09 E05
LOCKHEED	26	1914	N28 E44	SAC PEAK	28	1804	N08 W36	LOCKHEED	30	1546	N12 W57
LOCKHEED	26	2011	S17 W75	HAWAII	28	2012	N12 W41	LOCKHEED	30	1552	N10 W90
LOCKHEED	26	2041	N28 E43	HAWAII	28	2050	S12 E28	LOCKHEED	30	1820 E	N13 W50
LOCKHEED	26	2104	N22 W48	LOCKHEED	28	2216	N09 W12	LOCKHEED	30	1820 E	N13 W50
LOCKHEED	26	2119	S17 W75	LOCKHEED	28	2220	S08 E25	LOCKHEED	30	1830	N11 W68
HAWAII	26	2122 E	S15 W73	LOCKHEED	28	2240 E	N12 E38	LOCKHEED	30	1830	N11 W68
HAWAII	27	0026 E	N16 E10	HAWAII	29	0004 E	S09 E25	HAWAII	30	1906	N15 W50
LOCARNO	27	1336	N10 E56	LOCKHEED	29	1625	N30 E47	HAWAII	30	1934	N15 W50
SAC PEAK	27	1530	N10 E55	LOCKHEED	29	1702	N11 W52	LOCKHEED	30	2105 U	N11 W68
LOCKHEED	27	1658	N11 E27	LOCKHEED	29	1705	N26 W90	LOCKHEED	30	2149 D	N31 W09
LOCKHEED	27	1724	N18 W06	LOCKHEED	29	1726	S08 E16	MCMATH	31	1526 E	N10 W60
LOCKHEED	27	1739	N27 W57	LOCKHEED	29	1746	N10 W49	LOCKHEED	31	1540 E	S17 W71
LOCKHEED	27	1829	N14 W17	* SAC PEAK	29	1746	N10 W49	LOCKHEED	31	1550	N27 W20
LOCKHEED	27	1832	S18 W90	* HAWAII	29	1748 E	N07 W52	LOCKHEED	31	1555	N11 W61
HAWAII	27	1942	N20 E75	LOCKHEED	29	1755	N23 W90	LOCKHEED	31	1638	N28 W14
LOCKHEED	27	1946 E	N19 E73	LOCKHEED	29	1905	S07 E14	LOCKHEED	31	1713	N11 E01
HAWAII	27	2026	N08 W63	LOCKHEED	29	1938	N16 W88	SAC PEAK	31	1742	N12 W61
LOCKHEED	27	2211	N11 W27	LOCKHEED	29	1938	N16 W88	LOCKHEED	31	1742	N12 W59
HAWAII	27	2217 E	N10 W28	LOCKHEED	29	2020	N10 W53	HAWAII	31	1746	N11 W65
HAWAII	27	2230	N19 E67	HAWAII	29	2022	N20 W54	HAWAII	31	1846	N11 W90
LOCKHEED	27	2233 U	N21 E70	LOCKHEED	29	2045	S07 E13	LOCKHEED	31	1915	S12 E80
HAWAII	27	2254 E	N30 E29	LOCKHEED	29	2050	N10 W53	LOCKHEED	31	2128	N06 E90
ARCETRI	28	1003 E	N08 W26	HAWAII	29	2222	N10 W56	LOCKHEED	31	2133	N19 E07
CAPRI S	28	1415	N20 E57	HAWAII	29	2356	S10 E39	HAWAII	31	2134	S06 W14
								LOCKHEED	31	2351	N12 W67

*Rated as flare of importance ≥ 1 by other observatories (see CRPL-F 185 Part B).

COMMENCE - STANDARD - SOLAR

SOLAR FLARES

OCTOBER 1959

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION			DURA- TION — MINUTES	IN- FOR- TANCE	OBS. COND.	TIME — U T	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT
		START	END	APPROX.	MGMATH PLAGE REGION	MEAS. AREA Sq. Deg.					CORR. AREA Sq. Deg.	MAX. WIDTH Rg	MAX. INT. %	
				LAT.	MER. DIST.									
ONDREJOV	01	1121 E	1130	S17	E59	5401	9 D	1	3	1122			2.70	
	03	1008 E	1016	N08	W52	5389	8 D	1	1	1008	2.00			
{ ZURICH	04	1155 E	1210	S10	E28	5401	15 D	1	2	1155	2.00			
	04	1200 E	1214 D	S11	E28	5401	14 D	1	2	1200	1.80		71	
{ GOOD HOPE	04	2301	2320	S16	E09	5397	19	1	2	2303	1.90			
	04	2301	2320	S16	E09	5397	19	1	2	2303	1.90			
VOROSHILOV	05	1010 E	1020	S15	E08	5401	10 D	1	2	1010	1.00			
	05	1319 E	1323	S09	W51	5392	4 D	1	3	1319	2.00			
LOCARNO	05	1513 E	1520	S08	E00	5401	7 D	1	3	1513	2.00			
	05	1555	1606 D	N04	E40	5405	11 D	1	2	1600	1.00			
GOOD HOPE	06	0630 E	0643	N07	W67	5396	13 D	1	2	0630	3.30			
	06	0711	0732	N06	E28	5405	21	1	1	0716	3.60			
{ TASHKENT	06	0714	0719 D	N06	E30	5405	5 D	1	1	0716	6.06		85	
	06	0714 E	0721	N05	E28	5405	7 D	1	1	0718	1.82			
{ ABASTUMANI	06	1241 E	1248	S11	W68	5392	7 D	1	3	1241	2.00			
	06	1241 E	1316	N06	W69	5396	28	1	3	1254	1.00			
ZURICH	06	1329 E	1341	N30	E61	5408	12 D	1	3	1329	1.00			
	06	1343	1355	S11	W69	5392	12	1	2	1348	1.10			
GOOD HOPE	06	1405	1445	N32	E65	5408	40	1	2	1420	2.30			
	06	1406	1448	N30	E67	5408	42	2	2	1418	6.00			
{ ONDREJOV	06	1409	1445	N30	E62	5408	36	16	3	1420	2.30			
	06	1414	1437	N32	E60	5408	23	1	3	1418	6.00			S-SWF
MEUDON	06	1414	1437	N32	E60	5408	23	1	3	1418	6.00			
	06	1415 E	1445 D	N26	E62	5408	30 D	2	3	1430	2.00	5.80		
STOCKHOLM	06	1440	1500	N07	W71	5396	20	1	3	1447	1.20	3.70		
	06	1440	1500	N07	W71	5396	20	1	3	1447	1.20	3.70		
GOOD HOPE	06	2005	2037	N29	E61	5408	32	1	2	2013	2.00	4.20		
	06	2322	2344	N30	E62	5408	22	1	2	2334	2.35	4.90		
HUANCAYO	06	2322	2344	N30	E62	5408	22	1	2	2334	2.35	4.90		
	06	2359	0007	N30	E60	5408	8	1	2	0002	2.17	4.35		
VOROSHILOV	06	2359	0007	N30	E60	5408	8	1	2	0002	2.17	4.35		
	06	2359	0007	N30	E60	5408	8	1	2	0002	2.17	4.35		
KODAIKNI	07	0500 E	0510 D	N30	E55	5408	10 D	2	2	0500	2.20			
	07	0500 E	0510 D	N30	E55	5408	10 D	2	2	0505	1.70			
{ KODAIKNI	07	0500 E	0510 D	N30	E55	5408	10 D	2	2	0510	1.30			
	07	0502	0544	N32	E59	5408	42	16	2	0504	8.00			
TASHKENT	07	0502	0544	N32	E59	5408	42	16	2	0504	8.00			
	07	0505 E	0534	N30	E55	5408	29 D	2	2	0505	8.00			
{ SYDNEY	07	0505 E	0534	N30	E55	5408	29 D	2	2	0505	8.00			
	07	1033		N30	E55	5408	1	1	1					
{ UCCLE	07	1033		N30	E55	5408	1	1	1					
	07	1041	1048	N28	E51	5408	7	1	3	1043	2.10			
{ ONDREJOV	07	1103	1122 D	N10	W75	5394	9	1	3	1104	2.60			
	07	1103 E	1122 D	N10	W80	5394	19 D	1	3	1108	3.00			
{ CAPRI G	07	1155	1321 D	S18	W89	5392	86 D	1	3	1201	3.00			
	07	1210 E	1327 D	S18	W87	5392	77 D	1	3	1216	4.00			
{ GOOD HOPE	07	1300 E	1405	N32	E53	5408	65 D	16	3	1340	3.00			
	07	1305 E	1532 D	N30	E53	5408	147 D	1	3	1337	5.00			
LOCARNO	07	1328	1355	N30	E54	5408	27	16	3	1329	3.00			
	07	1426	1445	N29	E52	5408	19	2	3	1431	4.30			
{ ONDREJOV	07	1439 E	1457	N31	E51	5408	18 D	1	3	1439	5.00			
	07	1450	1540	N32	E53	5408	50	16	2	1510	4.00			
ZURICH	07	1458	1534	N29	E52	5408	36	16	3	1500	1.00			
	07	1529 E	1538	N08	E19	5405	9 D	1	3	1529	5.00			
{ ONDREJOV	07	1529 E	1545	N31	E50	5408	16 D	1	3	1529	1.00			
	07	1529 E	1545	N31	E50	5408	16 D	1	3	1529	5.00			
KYOTO	08	0415 E	0450 D	S10	W90	5392	35 D	1	3	0415	100			
	08	0415 E	0450 D	S10	W90	5392	35 D	1	3	0415	100			

SOLAR FLARES

OCTOBER 1959

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION			DURA- TION — MINUTES	IM- POR- TANCE	OBS. COND.	MEASUREMENTS				PROVISIONAL IONOSPHERIC EFFECT	
		START	END	APPROX. LAT.	MAGNIT. PHASE REGION	MEAS. AREA Sq. Deg.				CORR. AREA Sq. Deg.	MAX. WIDTH R _z	MAX. INT. %			
{ CAPRI S CAPRI G LOCARNO CAPRI S MEUDON ONDREJOV CAPRI G	08	0748	0807	S14	W28	5401	19	1	3	0759	2.20	2.60			
	08	1051 E	1137	N05	W05	5405	46 D	1	3	1104	5.00	5.00			
	08	1057	1114	N04	W04	5405	17	1	2						
	08	1425	1455	N03	W06	5405	30	2	2	1432	6.00	6.00			
	08	1428 E	1508 D	S01	W05	5405	40 D	2	2	1437	5.00	5.00			
	08	1430	1450	N05	W05	5405	20	1	3						
	08	1431 E	1441	N07	W07	5405	10 D	1	3	1437	8.00	8.00			
	08	1432 E	1452 D	N03	W02	5405	20 D	2	1	1442	9.00	9.00			
	09	1054	1104	N01	W17	5405	10	1	3	1059	2.30	2.62		45	
	09	1058 E	1107	N02	W18	5405	9 D	1	3	1058	3.00	3.00			
	09	1249	1304	N08	W08	5405	15	1	2	1254					
	{ ALMA-ATA SYDNEY TASHKENT ALMA-ATA SYDNEY SYDNEY LOCARNO ONDREJOV ZURICH LOCARNO GOOD HOPE ONDREJOV CAPRI G	10	0326 E	0546 D	S17	W37	5401	140 D	2	2	0409	3.95	9.50		
10		0329	0455	S17	W38	5401	86	2	1	0409	4.00	6.00			
10		0437 E	0624	S16	W49	5401	107 D	2	3	0506	16.52	27.00			
10		0456	0546 D	S19	W54	5401	50 D	2	2	0503	6.86	20.50			S-SWF
10		0457	0518	S10	W56	5401	21	2	1	0501	3.00	5.00			
10		0457	0523 D	S19	W54	5401	26 D	3	1	0505	11.00	21.00			
10		0836	0910	N31	E15	5408	34	1	3						
10		1029 E	1034	S08	E70	5418	5 D	1	3	1030	1.00	1.00			
10		1029 E	1034	S10	E74	5418	5 D	1	3	1029	2.00	2.00			
10		1029 E	1050	S16	W44	5401	21 D	1	1	1029					
10		1030 E	1035 D	S17	W44	5401	5 D	1	3						
10		1102	1114	S17	W57	5401	12	1	3	1104	1.10	2.10			
10	1242 E	1247	S17	W46	5401	12	1	3	1242	6.00	6.00				
10	1440 E	1455	N21	W20	5406	15 D	1	2	1442						
{ SYDNEY CAPRI G KIEV CAPRI S CAPRI G GOOD HOPE GOOD HOPE CAPRI G GOOD HOPE SYDNEY GOOD HOPE PIRCULI PIRCULI LOCARNO MEUDON PIRCULI MEUDON GOOD HOPE ZURICH ZURICH	11	0433	0454	S17	W54	5401	21	1	2	0436	1.50	3.00			
	11	0828	0834	S17	W58	5401	6	1	3	0829	3.00	3.00			
	11	1022	1032	N07	E85	5420	10	1	2	1024	.88	6.30			
	11	1123 E	1245 D	S16	W59	5401	22 D	1	2	1208	1.40	2.50		52	
	11	1147 E	1250 D	S17	W57	5401	63 D	1	1	1150	6.00	6.00			
	11	1148	1257	S17	W57	5401	69	1	2	1202	2.10	4.20			
	11	1358	1414	S18	W63	5401	69	1	1	1400	1.00	2.60			
	11	1400 E	1403 D	S17	W61	5401	3 D	1	2	1403	3.00	3.00			
	11	1442	1444 D	S04	E88	5418	2 D	1	2	1443	.50	6.00			
	11	2258	2304 D	S16	W66	5401	6 D	2	2	2259	2.00	6.00			
	12	0636 E	0653	N05	W50	5405	17 D	1	2	0636	1.80	2.80			
	12	0810	0815	N13	E01	5411	5	1	2	0813	.83	.90		44	
12	0843 E	0904 D	N05	W50	5405	21 D	1	2	0904	1.38	2.34		59		
12	0900 E	1030	N05	W51	5405	90 D	1	3							
12	0917	0948	N20	E00	5409	31	1	2	0945	2.75	3.00			48	
12	0922	0950	N30	W12	5408	28	1	2							
12	1234	1258	S20	E38	5416	24	1	1							
12	1234	1303	S20	E42	5416	29	1	1	1239	2.30	3.60				
12	1243 E	1249	S20	E39	5416	6 D	1	1	1243	1.00	1.00				
12	1243 E	1250	N03	W55	5405	7 D	1	1	1243	2.00	2.00				
12	1335 E	1349	N03	W55	5405	14 D	1	2	1335	1.00	1.00				
{ LOCARNO CAPRI G	13	0635	0645	N04	W63	5405	10	1	2						
	13	0713 E	0755	S16	W85	5401	42 D	2	3						

SOLAR FLARES

OCTOBER 1959

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION		DURA- TION — MINUTES	IM- POR- TANCE	OBS. COND.	MEASUREMENTS			PROVISIONAL IONOSPHERIC EFFECT	
		START	END	APPROX. LAT. MER. DIST.	MAGNITH PLAGE REGION				TIME U-T	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.		MAX. WIDTH He
{ GOOD HOPE GOOD HOPE CAPRI G TASHKENT SIMEIZ ONDREJOV LOCARNO LOCARNO ZURICH	13	0743 E	0805	S15 W88	5401	22 D	1	3	0.743	1.90			
	13	0815	0925	S19 W90	5401	70 D	2	3	0.821	1.20			
	13	0821 E	0835	S16 W85	5401	14 D	2	3					
	13	0827 E	0857 D	S19 W85	5401	30 D	16	3					
	13	0836 E	0838 D	N18 E88	5426	2 D	1	1		1.10	11.00	1.40	185
	13	0839 E	0907	S15 W85	5401	28 D	1	1		0.46	3.30	2.50	
	13	0845 E	0900	S06 E14	5424	15 D	1	2					
	13	0845 E	0910	S19 W80	5401	25 D	1	2					
	13	1358 E	1401	N00 W70	5405	3 D	1	2			1.00		
	14	0518 E	0549 D	S09 W39	5423	31 D	1	1		1.29	1.85		45
{ CAPRI G CAPRI G PIRCULI PIRCULI LOCARNO LOCARNO CAPRI G CAPRI S LOCARNO AROSA ZURICH	14	0558	0602	S08 E02	5424	4	1	1	0.600	0.73	0.82	45	
	14	1245 E	1252 D	N06 W85	5405	7 D	1	3	1.246	4.00			
	15	1139 E	1200 D	N28 W52	5408	21 D	1	3	1.149	1.19	2.13	45	
	15	1139 E	1200 D	S10 W57	5423	21 D	1	3	1.149	0.92	1.88	47	
	15	1230 E	1245 D	N29 W49	5408	15 D	1	1					
	16	0721 E	0726 D	S09 W04	5418	5 D	1	3	0.722		3.00		
	16	1023	1101 D	S05 W11	5418	38 D	16	3	1.036	2.60	2.00		
	16	1025	1120	S06 W09	5418	55	1	1	1.100				
	16	1040	1100 D	S05 W08	5418	20 D	1	1					
	16	1227 E	1232	N29 W63	5408	5 D	1	3	1.227		2.00		
{ SYDNEY PIRCULI ONDREJOV LOCARNO SIMEIZ GOOD HOPE CAPRI G AROSA PIRCULI ABASTUMANI	17	0052	0104	S08 W40	5424	12	1	2	0.057	1.50	2.00		
	17	0641 E	0735 D	N27 W76	5408	54 D	16	2	0.705	1.47	5.41	56	
	17	0751 E	0801 D	N31 W74	5408	10 D	1	3	0.754				
	17	0850	0920	S04 E53	5427	30	1	1	0.900		2.00		
	17	0850	0930 D	S06 E58	5427	40 D	1	1	0.855	0.90	1.60	66	
	17	0851	0915	S05 E56	5427	24	1	1	0.852	1.20	2.20		
	17	0851 E	0921	S05 E56	5427	30 D	1	3	0.853		3.00		
	17	0852	0900 D	S05 E54	5427	8 D	1	1					
	17	0853 E	0916	S04 E56	5427	23 D	16	2	0.858	1.10	3.18	64	
	17	1406 E	1408	S05 E57	5427	23 D	1	3	0.908	2.72	5.06	70	
{ VOROSHILOV VOROSHILOV VOROSHILOV TASHKENT ONDREJOV SIMEIZ CAPRI G VOROSHILOV ONDREJOV SIMEIZ CAPRI G KIEV CAPRI G ONDREJOV	17	2216 E	2320	N04 E01	5420	2 D	1	3	1.408	3.00	3.00		
	17	2337	0001	S07 E50	5427	64 D	16	3	2.218	3.34	5.19	73	
	17	2337	0001	N07 E46	5430	24	16	3	2.343	3.07	4.28	80	
	18	0112	0132	N29 W86	5408	20	1	2	0.121	0.81	4.77	78	
	18	0603	0617	N02 E59	5427	14	16	2	0.608	1.01	2.00	110	
	18	0610 E	0615	N02 E57	5427	5 D	1	3			3.80		
	18	0815 E	0830 D	N01 E58	5427	15 D	1	3	0.819	1.82	3.20	61	
	18	1117 E	1129	N02 E54	5427	12 D	1	3	1.117		3.00		
	18	2254	2258	N01 E49	5427	4	16	3	2.256	1.44	2.13	91	
	19	0624	0632 D	N02 E32	5427	8 D	1	1	0.627				
{ CAPRI G CAPRI G KIEV CAPRI G CAPRI G ONDREJOV	19	0706	0715	S03 E31	5427	9	1	2	0.707	0.90	1.00	74	
	19	0708 E	0720	S03 E30	5427	12 D	1	3	0.710		4.00		
	19	0807 E	0842 D	N09 W22	5420	35 D	1	3	0.808		3.00		
	19	1023	1027	S05 W50	5418	4	1	2	1.025		4.41	66	
	19	1316 E	1356	S04 E11	5425	40 D	1	3	1.322	3.11	4.41		
	19	1320	1420	S05 E27	5427	60	1	3	1.322		4.00		
	19	1322	1341	S05 E14	5425	19	1	3	1.329		4.00		
													S-SWF S-SWF

SOLAR FLARES

OCTOBER 1959

OBSERVATORY	DATE	OBSERVED TIME		LOCATION			DUR. TION MINUTES	IM. POR. TANCE	OBS. COND.	MEASUREMENTS				PROVISIONAL IONOSPHERIC EFFECT	
		START	END	APPROX. LAT.	MER. DIST.	MAGN. PLAGE REGION				TIME U T	MEAS. AREA Sq. Deg.	CORR. AREA Sq. Deg.	MAX. WIDTH H _z		MAX. INT. %
{ ONDRE JOV CAPRI G CAPRI S CAPRI G KYOTO	19	1340	1354	S06 E26	5427	14	1	3	1351			1.90			
	19	1356	1414 D	N08 E27	5430	18 D	1	3	1400			2.30			
	19	1357 E	1430	N09 E21	5430	33 D	2	3	1405		8.00				
	19	1358 E	1420	N17 E22	5428	22 D	1	3	1407		2.50				
	19	1507 E	1530 D	N08 E90	5433	23 D	1	2	2335		1.45	1.16	120		
	19	2335 E	2344 D	S07 W55	5418	9 D	1								
	GOOD HOPE	20	0630 E	0656	N05 E85	5433	26 D	1	2	0630	1.00				
		20	0837 E	0910	N07 E16	5430	33 D	1	2	0853	.90	1.00		63	
		20	0843	0849 D	N08 E81	5433	6 D	1	2	0844	.46	2.10		66	
		20	0845 E	0848	N06 E78	5433	3 D	1	3	0846		4.00			
20		1210 E	1245 D	N06 E75	5433	35 D	1	3	1215		5.00				
20		1237 E	1245 D	N06 E08	5430	8 D	1	3	1239		3.00				
20		1326	1412	S04 E15	5427	46	1	3	1333	3.20	3.40				
20		1330 E	1402 D	S03 E14	5427	32 D	1	3	1333		5.00				
20		1416 E	1435 D	S05 W01	5425	19 D	1	3	1417		3.00				
21		0756	0810	S10 W74	5418	14	1	2	0805	1.08	3.40	1.60	63		
{ KRASNAYA ABASTUMANI PIRCULI	21	0758	0805	S10 W73	5418	7	1	2	0800	1.82	2.70		70		
	21	0802 E	0810 D	S09 W71	5418	8 D	1	3	0803	1.09	3.94		76		
	21	0806 E	0821	N06 E69	5433	15 D	1	3	0814	1.01	1.40		52		
	21	0834 E	0845	S05 E03	5427	11 D	1	3	0835		2.30				
	21	0836	0905	S04 E03	5427	29	16	3	0848	2.02	2.20		56		
	21	0952	1006	S02 W75	5418	14	16	2	0956	1.82	2.70		100		
	21	1000	1006	S06 W70	5418	6	1	3	1003	.64	1.99		46		
	21	1139 E	1145 D	N05 E67	5433	6 D	1	3	1140		5.00				
	21	1142	1155 D	N06 E68	5433	13 D	1	3	1144			2.90			
	22	0658 E	0712 D	S11 E90	5438	14 D	1	3	0658	2.08		1.36	140		
KYOTO	22	0709 E	0715	S11 W17	5431	6 D	1	3	0710						
	22	0758 E	0821 D	S10 E89	5438	23 D	1	2	0801	1.46	4.00				
	22	1020 E	1035 D	S09 E77	5438	15 D	1	3	1021		4.00				
	22	1236	1259	N13 W66	5421	23	1	3	1241	1.20	2.80				
	22	1248 E	1257 D	S11 W17	5431	9 D	1	3	1249		4.00				
	22	1424	1455 D	N04 E48	5433	31 D	1	3	1426		2.00				
	SYDNEY	23	0302	0327 D	S11 W27	5431	25 D	1	2	0313	3.00	3.00			
		23	0729	0740	N04 W13	5427	11	1	3	0734	.83	.93		45	
		23	0738	0755	S13 W13	5431	17	1	3	0744	1.10	1.28		56	
		23	0740	0800	N06 E36	5433	20	1	3	0750	.75	.75		58	
23		1120 E	1132 D	N07 E40	5433	12 D	1	3	1123		5.00				
23		1502	1520	N05 E30	5433	18	1	3	1515		2.00				
23		1517	1530 D	S08 W33	5431	13 D	26	3	1525		9.00				
23		1517	1532 D	S09 W34	5431	15 D	2	2	1520		7.00				
24		0747	0814 D	S05 W38	5431	27 D	1	3	0802	1.80	2.30				
24		0750 E	0756 D	S08 W39	5431	6 D	1	3	0752		3.00				
{ GOOD HOPE PIRCULI	24	0754	0815	S12 W41	5431	11	1	3	0755	.81	1.12		45		
	24	0759	0810	S03 W39	5427	21	1	3	0758	1.60	2.10				
	24	0822	0837	S02 W39	5427	11	1	2	0804	1.01	1.43		52		
	24	0938 E	0958	N04 E21	5433	15	1	2	0828	.46	.53		56		
	24	0938 E	0958	S12 W30	5431	20 D	1	3	0940		3.00				
	25	0128	0234 D	N40 W50	5428	66 D	2	3	0150	6.00	11.00				

SOLAR FLARES

OCTOBER 1959

OBSERVATORY	DATE	OBSERVED UNIVERSAL TIME		LOCATION			DURA- TION — MINUTES	IM- POR- TANCE	OBS. COND.	MEASUREMENTS				PROVISIONAL IONOSPHERIC EFFECT
		START	END	APPROX.	MATH	MEAS.				CORR.	MAX.	MAX.		
				LAT.	MER. DIST.	PLAGE REGION				AREA Sq. Deg.	AREA Sq. Deg.	WIDTH Rc	INT. %	
PIRCULI -LOCARNO CAPRI G LOCARNO LOCARNO	25	0700	0710	N13	E11	5433	10	1	1	1.19	1.31		48	
	25	1025	1055	S09	E44	5438	30	3	3					
	25	1200 E	1212	N04	E08	5433	12 D	3	3	4.00				
	25	1433	1440	S12	E42	5438	7	3	3					
CAPRI G ONDREJOV	25	1452	1505	N09	E22	5437	13	3	3	1.00				
	26	0745	0758 D	N05	E02	5433	13 D	3	3	4.00		1.90		
GOOD HOPE CAPRI G GOOD HOPE GOOD HOPE	26	1045 E	1101	N05	E01	5433	16 D	3	3					
	27	0739	0750	S16	W63	5431	11	1	1	1.20	2.80			
	27	0947 E	0952 D	N05	W13	5433	5 D	3	3	4.00				
	27	1034	1052	S15	W77	5431	18	1	1	.90	0948			
KRASNAYA	27	1050	1102	N09	W15	5433	12	1	1	2.40	2.50			
	28	1045	1106	N29	E30	5440	21	1	1	2.27	1.40		100	
{ SYDNEY KYOTO KYOTO KRASNAYA	31	0155	0216	S10	W34	5438	21	1	2	2.00	2.00		100	
	31	0158 E	0205 D	S10	W28	5438	7 D	1	1	1.25			100	
	31	0158 E	0207 D	S08	W33	5438	9 D	1	1	3.74			100	
	31	0914	0919	N11	W31	5437	5	2	2	1.40	.80		115	


COMMERCE - STANDARDS - BOULDER

CAPRI G CAPRI S GOOD HOPE KIEV* KODAIKANAL KODAIKANAL KRASNAYA LOCKHEED	AMACAPRI - GERMAN ANACAPRI - SWEDISH ROYAL OBSERVATORY, CAPE OF GOOD HOPE KIEV UNIVERSITY KODAIKANAL KRASNAYA PAKHRA LOS ANGELES	MOSCOW - G AISH ROYAL OBSERVATORY, EDINBURGH HERSTMONCEUX SAC PEAK SACRAMENTO PEAK SCHAUTINSLAND UNITED STATES NAVAL RESEARCH LABORATORY USNRL
--	--	---

SAC PEAK: ALL VALUES IN MAX. INT. COLUMN ARE ARBITRARY UNITS (0-40), NOT PERCENT OF CONTINUOUS SPECTRUM.
 E - LESS THAN & - PLUS
 D - GREATER THAN - - MINUS
 U - APPROXIMATE □ - NOT REPORTED

LOCKHEED OBSERVATIONS: ALL VALUES IN THE MAXIMUM INTENSITY COLUMN ARE ARBITRARY UNITS ON A SCALE OF 1 TO 4 - NOT PERCENT OF THE CONTINUOUS SPECTRUM.

INTERVALS OF NO FLARE PATROL OBSERVATIONS
OCTOBER 1959



Stations Include:

COMMERCE - STANDARDS - BOULDER 202041-102-10

- | | | | |
|--------------------|-----------------|-----------------------------|-----------------|
| Abastumani | Hawaii | McMath | Sacramento Peak |
| Alma Ata | Huancayo | Meudon | Simeiz |
| Anacapri (Swedish) | Kharkov | Mitaka | Sydney |
| Arcetri | Kiev GAO | Moscow Gaish | Tashkent |
| Arosa | Kodaikanal | Nizamia | Uccle |
| Athens | Krasnaya Pakhra | Ondrejov | Utrecht |
| Climax | Kyoto | Pirculi | Voroshilov |
| Dunsink | Locarno | Royal Greenwich Observatory | Zurich |
| Good Hope | Logkheed | Herstmonceux | |

IONOSPHERIC EFFECTS OF SOLAR FLARES

(SHORT-WAVE RADIO FADEOUTS)

DECEMBER 1959

Dec. 1959	Start UT	End UT	Type	Wide Spread Index	Importance	Observation Stations	Known Flare, UT CRPL-F 185
1	1247	1327	S-SWF	1	3	JU	1208E
1	1512	1613	Slow S-SWF	5	2+	BE, FM, HU, LA, MC, NE, PR, WS, CW* *	1456E
1	1705	1900	S-SWF	5	3	AN, BE, FM, HU, LA, MC, NE, PR, WS, CW***	1638
2	0330	0450	S-SWF	1	3-	OK	*
2	0500	0550	Slow S-SWF	1	2	OK	*
2	1246	1402	S-SWF	5	2+	BE, DA, HU, MC, NE, PR, PU, SW, CW***	1219E
3	1017	1047	S-SWF	1	2	NE	1010
3	1414	1500	S-SWF	3	2-	HU, PR	1408E
3	1757	1903	S-SWF	5	2+	AN, BE, FM, HU, LA, MC, NE, PR, WS, CW*	1756
4	0028	0208	S-SWF	5	2+	AD, OK	0032
4	0210	0240	G-SWF	1	1+	OK	0208
4	0800	0840	-	1	-	CW**	*
4	1820	1920	S-SWF	5	2+	BE, FM, HU, LA, MC, PR, WS	1814
5	0620	0633	S-SWF	1	3	KO	*
5	1003	1022	S-SWF	3	2	NE, PU	1004E
5	1220	1232	S-SWF	5	2	BE, NE, PR, PU	1230E
5	1615	1630	S-SWF	5	1	BE, MC, PR, WS	
7	0440	0520	S-SWF	5	2-	CA, KE, OK	0434
7	1042	1107	S-SWF	5	1	PR, PU	*
8	0118	0200	S-SWF	5	1+	AD, CA, OK	0120E
8	0755	0820	S-SWF	5	1+	CA, KO, OK, NE	*
10	0518	0545	S-SWF	5	1+	CA, KO, OK	*
10	2355	0037	Slow S-SWF	5	2	AD, CA, OK	*
11	0407	0430	S-SWF	1	1+	OK	*
12	0800	0810	-	1	-	CW++	0812
17	0400	0422	Slow S-SWF	4	1+	OK, TO	*
18	0637	0657	S-SWF	4	1+	KO, OK	*
19	0345	0405	S-SWF	1	1	OK	*
24	0343	0450	Slow S-SWF	1	2	OK	*

CA = Canberra, Australia
 DA = Darmstadt, G.F.R.
 JU = Juhiesruh, G.D.R.
 KE = Kerguelen
 KO = Kodaikanal, India
 LA = Los Angeles, Calif.
 NE = Nederhorst den Berg, Netherlands

PU = Prague, Czechoslovakia
 TO = Hiraizo Radio Wave Observatory, Japan
 SW = Enkoping, Sweden
 CW* = Cable and Wireless, Barbados
 CW** = Cable and Wireless, Somerton, England
 CW*** = Cable and Wireless, Brentwood, England
 CW++ = Cable and Wireless, Singapore

IONOSPHERIC EFFECTS OF SOLAR FLARES

(Sudden Cosmic Noise Absorption
Sudden Enhancements Of Atmospherics)
Solar Noise Bursts At 18 Mc.
SEPTEMBER 1959

SEPT. 1959	CLASS			WIDESPREAD INDEX	TIME (UNIVERSAL TIME)			PERCENT ABSORPTION SCNA	OBSERVATION STATIONS
	SCNA	SEA	Burst		BEGIN	MAX.	END		
1	1			1	0206	0211	0225	15	HA
1			1+	3	1420	1425	1428		MC, RE
1		2		5	1655	1708	1735		BO, DU, HA, NE, PA, SP
1	3			5	1656	1701	1800U	80	BO, HA, MC, RE, SP
1		2		5	1730		2355		BO, HA, MC, SP, (Noise storm, peaks at 1810-1830 (RE) 1950- 2007 (RE) 2228-2307)
1		2+		3	1928	1940	2007U		A1, A3, A5
1		2		3	2007	2022	2100		A1, A3, A5
2		1		3	1311		1329		JU, KU
2			1	5	1605		1607		BO, MC, RE, SP
2		1+		5	1606	1615	1640		A3, A5, BO, DU, NE, PA, SP
2	2			5	1607	1610	1635	55	BO, MC, RE, SP
2	1			4	1730	1747	1810	15	BO, MC
2		1		1	1735	1752	1810		BO
2			1	5	1800		0026		BO, HA, MC, (Noise Storm)
3			2	1	1425		1605		MC, (Group of bursts)
3			2	4	1739		1900		BO, MC, SP (Group of bursts)
3			1	1	2115		0030		HA (Noise Storm)
4		1+		3	1605	1632	1710		A2, A5
4			1	5	1700		2345		BO, HA, (Noise Storm)
4			2	1	2320		2326		HA
5			2	5	1554		1600		BO, MC, RE, SP
6			1	1	2045		2048		HA
6			2	1	2158		2205		HA
7			1	1	1615		1623		MC
9		2		1	0703		0741		NE
9	1			5	1557	1605	1625	20	BO, MC, RE, SP
9		2		5	1558	1611	-		BO, MC, NE, SP
9			1	5	1620		1625		BO, MC, RE, SP
9			1	5	1655		1658		BO, MC, SP
9			1	4	1711		1713		BO, MC
9			1	4	1824		1827		BO, MC
9			1	5	1833		1839		BO, MC, SP
9			1	1	2235		2240		MC
11			1	5	2000		2001		BO, HA, MC
12			1	5	2038		2050		BO, HA, MC, SP
13	1			5	1725	1740	1800	10	BO, HA, MC
14		1		1	0745		0821		NE
14			1	5	1833		1835		BO, MC, SP
14	1			5	2156	2200	2210	25	BO, HA, SP
15			1	5	2107		2109		BO, HA
16	1			4	1846	1850	1902	15	BO, MC
16		1		5	1846		-		BO, HA, MC
19			1	5	1939		1948		BO, HA, SP, (Group of bursts)
19			1	5	2027		2030		BO, HA, MC, SP
19			1	1	2203		2208		HA
19			1	1	2211		2218		HA
20			1	5	2147		2148		BO, HA, MC
20	1			1	2255	2300	2312	10	HA
21			1	1	0121		0130		HA
22			1	4	1652		1658		BO, MC
23			1	1	1909	1915	1917		RE
25			1	1	1444		1451		MC
25			2	5	1609		1617		BO, MC, RE
25			1	4	1625		1635		BO, MC
26			1	5	1830		2330		BO, HA, MC, SP, (Noise Storm)
29			1	5	2012		2015		BO, HA
29			1	5	2031		2035		BO, HA
29			1	1	2103		2105		HA

SOLAR RADIO EMISSION OUTSTANDING OCCURRENCES

Ottawa

JANUARY 1960

2800 Mc

Jan. 1960	Type	Start UT	Duration Hrs:Mins	Maximum		Remarks
				Time UT	Peak Flux	
3	2 Simple 2	1725	2	1725.7	12	
8	3 Simple 3	1805	2 30	1900	15	
10	1 Simple 1	1617.5	2	1618	7	
11	2 Simple 2	2056	>35	2108	220	In sunset oscillations
12	6 Complex f	1647.3	9	1649	80	
13	6 Complex f	1446	7	1449.5	18	
	4 Post Increase		15		6	
13	2 Simple 2	1847	4	1847.8	30	
15	6 Complex f	b1340*	>1 40	1357	700	*In interference
	4 Post Increase A		2 30		25	
	2 Simple 2 f	1730.5	8	1732	300	
16	3 Simple 3	b1543	>2' 17	1620	10	
17	3 Simple 3 A	b1600	>3 30	indet.	12	
	3 Simple 3	1616	40	1619	8	
19	3 Simple 3 A	1925	>2 10	2007	20	
	6 Complex	1936.5	25	1945	65	
22	2 Simple 2	1649	1.5	1649.3	10	
23	1 Simple 1	1519.5	2	1520.2	6	
24	1 Simple 1	1633	1.5	1633.7	6	
24	1 Simple 1	1715.3	1	1715.5	7	
25	2 Simple 2	1712.5	3.5	1714.2	90	
	4 Post Increase		1 15		6	
30	3 Simple 3 f	2015	12	2021	10	

CGMG/RC - STANDARDS - BOULDER

SOLAR RADIO EMISSION
OUTSTANDING OCCURRENCES
JANUARY 1960

IVb

BOULDER

167 MC

Jan. 1960	Type	Start UT	Time of Maximum UT	Duration Minutes	Intensity
3	3	1607.1	1607.2	1.2	1
3	3	1923.4	1923.4	0.1	1
3	3	2133.0	2133.0	0.2	1
3	3	2225.4	2225.6	0.3	2
3	3	2258.3	2258.8	0.7	1**
3	3	2320.5	2320.5	0.1	1**
3	3	2325.8	2325.8	0.2	1**
4	6	1423 E		165 D	2
4	3	1806.9	1806.9	0.1	1
4	7	1906		97	2
5	3	1916.5	1916.5	0.1	2
5	3	1919.6	1919.6	0.2	2
6	2	1500.9	1502.8	4.1	1*
6	2	2319	2320	12	1**
7	3	2224.0	2224.0	0.1	1
7	3	2302.3	2302.3	0.1	1**
7	3	2305.4	2305.4	0.2	2**
7	3	2307.6	2308.3	1.2	2**
8	6	1423 E		554 D	1
9	6	1422 E		458 D	1
10	3	1836.0	1837.1	2.0	2
10	3	1842.3	1842.3	0.3	2
10	3	2021.5	2021.5	0.1	1
10	3	2143.7	2143.7	0.2	2
10	7	2229		71 D	2
11	6	1421 E		562 D	2
11	9	2056.0		167 D	3**
12	6	1422 E	1935 U	561 D	2
12	8	1648.9	1651.2	12	3
13	6	1421 E	1957 U	562 D	1
14	6	1421 E	1756 U	564 D	2
15	6	1418 E		567 D	1
16	6	1422 E		565 D	2
16	8	2247	2250.0	6	3
17	6	1421 E		566 D	1
17	2	1910.0	1911.0	1.6	2
18	2	1705.0	1705.6	1.0	2
18	2	1739	1740	7	2
18	3	1757.0	1757.0	0.2	2
18	3	1912.8	1912.8	0.3	2
18	3	2330.2	2330.2	0.2	2**
19	3	1431.6	1431.7	0.8	2*
19	3	1433.8	1433.8	0.1	2*
19	3	1455.8	1455.8	0.3	2
19	3	1640.0	1640.0	0.1	1

Jan. 1960	Type	Start UT	Time of Maximum UT	Duration Minutes	Intensity
19	7	1849		300 D	1
19	9A	1941.0	1946.2	12	2
19	9B	1953	1956.5	8	1
20	2	1432.0	1433.1	2.2	2*
20	3	1802.0	1803.0	1.2	2
20	3	1813.0	1813.0	0.1	1
21	3	1458.0	1458.0	0.1	1*
21	3	1500.2	1500.2	0.1	1*
21	3	1602.0	1602.0	0.2	1
21	3	1739.9	1739.9	0.1	1
21	3	2337.7	2337.7	0.2	1**
22	3	1423.0	1423.0	0.1	1*
22	3	1550.5	1550.5	0.2	2
22	3	1805.4	1805.4	0.2	2
24	3	2003.5	2003.5	0.3	2
26	3	1422.0	1422.2	0.1	1*
26	3	1425.0	1425.0	0.2	1*
26	3	1507.2	1507.2	0.4	1
26	3	1643.9	1643.9	0.6	2
27	2	1417.8	1417.8	3.2	3*
27	3	1428.0	1428.0	0.2	2*
27	3	1504.7	1504.7	0.5	2
27	3	1531.3	1531.3	0.4	1
27	3	1543.0	1543.0	0.8	1
27	3	1805.5	1806.0	0.7	2
27	3	1809.0	1809.0	0.2	2
27	3	1838.5	1839.0	1.2	1
27	3	1852.0	1852.5	1.0	1
27	3	2348.8	2348.9	0.8	2**
28	3	1432.5	1432.5	0.5	2*
28	3	2334.5	2334.5	0.3	2**
29	3	1823.2	1823.2	0.2	2
30	3	1546.0	1546.0	1.0	2
30	3	1640.0	1640.3	0.4	2
30	2	1714	1714.0	7	2
30	3	1723.0	1723.0	0.5	3
30	3	1836.0	1836.0	0.9	1
30	3	2116.6	2116.6	0.4	2
30	3	2201.0	2201.0	0.2	3
30	3	2348.6	2348.6	0.3	1**
31	3	1423.0	1423.0	0.2	1*
31	3	1451.2	1451.2	0.1	2*
31	3	2002.0	2002.0	0.8	2
31	3	2350.9	2351.2	1.6	2**
31	3	2354.1	2355.2	2.8	2**

* On sunrise pattern

** On sunset pattern

TIMES OF OBSERVATION


Jan. 1960	U.T.	Jan. 1960	
1	1422-2329	16	1422-2347
2	1424-2333	17	1421-2347
3	1424-2334	18	1645-2348
4	1423-2335	19	1419-2349
5	1423-2335	20	1419-2350 I after 1830
6	1422-2337	21	1417-2352 I 1915-2030
7	1423-2337	22	1415-2354
8	1423-2337	23	1416-2356
9	1422-2200	24	1416-2356
10	1730-2340	25	1415-2357 I throughout day
11	1421-2343	26	1415-0000 I throughout day
12	1422-2047	27	1414-0000
	2103-2343	28	1415-0002
13	1421-2343	29	1410-0004 I 1830-0004
14	1421-2345	30	1420-0005 I 1725-1745
15	1418-2345	31	1410-0006

SOLAR RADIO EMISSION
INTERFEROMETRIC OBSERVATIONS

JANUARY 1960


Nancay

169 Mc



No observation on the 16th


JANUARY 1960



GEOMAGNETIC ACTIVITY INDICES

DECEMBER 1959

Dec. 1959	C	Values Kp								Sum	Ap	Final Selected Days	
		Three hour Gr. interval											
		1	2	3	4	5	6	7	8				
1	1.1	5o	4+	4+	5-	4+	3+	3o	2-	31-	28	Five Quiet	
2	1.2	3+	4+	4+	5o	4o	4+	4o	4o	33+	30		
3	1.5	5o	5-	5o	5+	5+	6-	4o	5+	40+	50		
4	0.7	3+	4o	3-	3-	2o	2-	2+	2+	21o	12		7
5	1.7	0o	0o	4o	5-	6+	7-	8-	6+	36-	68		10
6	0.9	3o	4+	3+	3+	3o	2o	3o	2+	24+	16	11	
7	0.3	3o	3-	1-	0+	2-	2-	1-	1+	12o	6	21	
8	0.3	1+	2o	2-	1o	1-	3-	2+	2o	14-	7	22	
9	0.3	1+	1+	3-	3-	3o	2o	1-	2o	16-	8		
10	0.2	2+	2-	2-	1-	0+	1+	2+	1+	12-	6		
11	0.2	0+	2o	2-	2o	1+	2+	1-	2-	12o	6	Five Disturbed	
12	1.0	2+	4+	3-	3o	3-	2o	3o	4o	24o	16		
13	1.1	4o	3+	2+	2-	1o	2+	3o	5+	23o	18		
14	1.4	5-	4+	5+	5o	5-	5+	5-	3o	37o	40		3
15	1.1	3+	4+	4+	4o	3+	3+	4o	4+	31o	26		5
16	1.0	3+	2+	3o	3+	4-	4o	2+	3o	25o	17	14	
17	0.3	3o	1+	3-	3-	2o	1-	1+	2-	15+	8	27	
18	0.5	2o	2-	3o	3-	4-	2o	1+	2o	18+	10	28	
19	0.8	2+	3-	3+	4-	4o	3-	2o	1o	22-	14		
20	0.3	2+	2o	1+	2-	1+	2o	2-	2-	14o	6		
21	0.1	1+	1+	1+	1-	1-	0+	1o	1o	8-	4	Ten Quiet	
22	0.4	2-	0+	0+	0+	1o	2+	2o	3o	11o	6		
23	1.3	3-	3-	4-	3-	3-	6o	4+	5-	29+	28		7
24	0.9	4o	1+	4o	4o	3+	1+	2-	2o	22-	15		8
25	0.4	2o	3o	2o	3-	2-	2-	2+	1+	17-	8		9
26	1.2	3-	4-	3+	4+	4+	4-	4o	5-	31-	26	10	
27	1.4	5-	5-	4o	6-	4+	5-	4o	4+	36+	38	11	
28	1.4	5o	5-	4+	5-	5+	5+	5-	3o	37o	40	17	
29	0.9	4o	4o	4o	4-	4-	3-	1o	2o	25o	18	20	
30	0.8	2-	3o	3o	4-	4-	3o	3-	1o	22-	14	21	
31	0.4	2-	2-	3-	2o	1o	1o	2+	1+	14-	7	22	
31	0.4	2-	2-	3-	2o	1o	1o	2+	1+	14-	7	31	
Mean:	0.81									Mean:	19		



CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS
NORTH ATLANTIC

DECEMBER 1959

Dec. 1959	North Atlantic 6-hourly quality figures				Short-term forecasts issued about one hour in advance of:				Whole day index	Advance forecasts (J-reports) for whole day; issued in advance by:				Geomag- netic K _{Pr}	
	00 to 06	06 to 12	12 to 18	18 to 24	00	06	12	18		1-7 days Final	1-7 days Js	1-7 days SDW	1-7 days J	Half Day (1) (2)	
1	5o	5-	6+	5o	4	4	6	4	5+	4		4	(4)	3	
2	5+	5o	6-	4o	5	4	6	5	5-	5		5	(4)	(4)	
3	5-	4+	6-	4+	4	4	6	5	5-	4		4	(5)	(4)	
4	5-	5o	7o	6+	4	4	6	6	6-	5		5	3	1	
5	6+	6-	6o	5-	5	6	6	6	6-	5		5	2	(5)	
6	5+	5o	7o	6+	4	4	5	6	6o	6		6	3	2	
7	6-	6o	7o	7-	5	5	7	7	6+	7		7	1	1	
8	6+	6+	7+	6-	6	6	7	7	6+	7		7	1	2	
9	5+	6+	7o	7o	6	6	7	6	6+	7		7	2	2	
10	7-	6o	7o	6+	6	6	7	7	6+	6		6	1	1	
11	6+	6-	7-	7-	7	5	7	7	6+	6		6	1	1	
12	6o	6+	7o	6o	6	5	7	7	6+	7		7	3	3	
13	5-	6o	7+	5+	6	5	7	6	6o	7		7	3	3	
14	4+	5-	6o	5+	5	4	6	6	5o	7		7	3	(4)	
15	5+	4+	6+	4+	5	5	6	6	5o	6		6	(4)	3	
16	5o	5+	6o	4+	4	4	6	5	5o	6		6	3	2	
17	5o	6-	7+	5-	5	6	7	6	6-	6		6	2	1	
18	5o	6-	7o	7-	5	5	7	6	6o	6		6	2	2	
19	5+	5+	7-	6+	6	5	7	7	6-	6		6	3	1	
20	5o	5+	7+	6+	5	5	7	7	6o	5		5	2	1	
21	6-	6o	7o	7-	5	6	7	7	6+	5		5	1	1	
22	6+	6o	7o	6-	6	6	7	6	6+	6		6	0	1	
23	5-	5o	7o	4+	5	5	7	5	5+	6		6	2	(4)	
24	5-	5o	7-	6o	4	5	7	5	6-	4		4	3	2	
25	5o	6-	7-	6+	6	5	7	6	6o	5		5	2	1	
26	6-	6-	7-	5o	5	5	7	6	6-	6		6	(4)	3	
27	5o	5o	6+	4+	5	5	6	5	5o	6		6	(4)	(4)	
28	5-	4+	6-	5-	5	5	6	5	5-	6		6	(4)	(4)	
29	5o	5o	6+	6-	5	5	6	5	6-	6		6	(4)	2	
30	5o	6-	6-	5o	6	5	7	6	5+	6		6	3	2	
31	5-	5+	7-	6-	5	5	6	6	6-	6		6	2	1	
Score: Quiet Periods					P	13	14	27	9		11		11		
					S	17	14	3	16		18		18		
					U	0	0	1	0		1		1		
					F	0	0	0	0		1		1		
Disturbed Periods					P	0	1	0	0		0		0		
					S	1	2	0	5		0		0		
					U	0	0	0	0		0		0		
					F	0	0	0	1		0		0		

() represent disturbed values.


CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS NORTH ATLANTIC

VIb

— Short-term forecast
○ Quality figure


DECEMBER 1959

| Range of reports





OUTCOME OF ADVANCED FORECASTS



FINAL ESTIMATE



USEFUL FREQUENCY RANGES -- NORTH ATLANTIC PATH

DECEMBER 1959





CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS

NORTH PACIFIC

DECEMBER 1959

Dec. 1959	North Pacific 12-hourly quality figures		Short-term fore- casts issued at		Whole day index	Advance forecasts (Jp reports) for whole day; issued in advance by:				Geomag- netic K _{SI}	
	0700 to 1900	1900 to 0700	0600	1800		1-7 days Final	1-7 days Jps	1-7 days SDW	1-7 days Jp	Half Day (1) (2)	
1	6	6	4	5	6	4		4	(5)	(4)	
2	6	6	4	5	5	5		5	(4)	(4)	
3	4	4	5	5	(4)	5		5	(5)	(6)	
4	6	5	5	5	5	5		5	3	2	
5	3	2	5	5	(2)	6		6	2	(8)	
6	6	6	5	6	6	6		6	3	2	
7	5	5	6	5	5	6		6	1	2	
8	5	6	5	5	5	6		6	1	1	
9	6	5	5	6	6	6		6	2	1	
10	5	5	5	5	5	5		5	1	1	
11	5	4	5	5	5	5		5	1	1	
12	7	5	5	5	6	6		6	2	2	
13	6	6	5	5	6	6		6	2	2	
14	5	5	5	5	(4)	6		6	(5)	(4)	
15	5	5	4	5	5	6		6	(4)	(4)	
16	5	5	5	5	5	5		5	3	(4)	
17	6	6	5	6	6	5		5	2	2	
18	7	7	5	6	7	5		5	2	2	
19	6	6	5	6	6	6		6	(4)	3	
20	6	5	5	6	5	5		5	1	2	
21	6	6	5	5	6	5		5	0	0	
22	5	6	5	5	6	6		6	0	2	
23	5	6	5	5	6	6		6	2	(4)	
24	6	5	5	6	6	5		5	(4)	2	
25	6	7	5	6	6	6		6	2	2	
26	5	6	6	5	6	6		6	(4)	(4)	
27	5	5	5	5	5	5		5	(4)	(4)	
28	4	5	4	4	(4)	5		5	(4)	(5)	
29	5	5	5	5	5	5		5	(4)	3	
30	6	6	5	6	6	5		5	2	3	
31	6	6	6	6	6	6		6	1	2	
Score:	Quiet Periods		P 10	14		18					
			S 14	14		7					
			U 2	0		1					
			F 2	0		1					
	Disturbed Periods		P 1	0		0					
			S 1	2		2					
			U 1	0		0					
			F 0	1		2					

() represent disturbed values.

COMMERCE - STANDARDS - BOULDER

Errata: In CRPL-F 185 Part B the score for December 1959 under "Whole day index" should have been under "1-7 days Final".


CRPL RADIO PROPAGATION QUALITY FIGURES AND FORECASTS

Vif

NORTH PACIFIC

DECEMBER 1959

OUTCOME OF ADVANCED FORECASTS FINAL ESTIMATE



ALERT PERIODS AND SPECIAL WORLD INTERVALS

INTERNATIONAL WORLD DAY SERVICE
JANUARY 1960

Issued Day/Time UT Jan 1960	Advance Geophysical Alert	No.	World-Wide Geophysical Alert	Special World Interval
11/1600		44	Magnetic Storm 10/0719Z	
12/0000	Burbank Solar Flare 11/2140Z			
14/1005	Fort Belvoir Magnetic Storm 13/1900Z			
14/1600		45	Magnetic Storm 13/1900Z	
18/1115	Fort Belvoir Magnetic Storm 17/12XXZ			
18/1600		46	Magnetic Storm 17/1200Z	
21/1600		47	Magnetic Storm 21/00XXZ	

COMMERCE - STANDARDS - BOULDER